The Checkerboard Copula and Dependence Concepts*

Liyuan Lin[†], Ruodu Wang[†], Ruixun Zhang[‡], and Chaoyi Zhao[§]

Abstract. We study the problem of choosing the copula when the marginal distributions of a random vector are not all continuous. Inspired by four motivating examples, including simulation from copulas, stress scenarios, corisk measures, and dependence measures, we propose to use the checkerboard copula, that is, intuitively, the unique copula with a distribution that is as uniform as possible within regions of flexibility. We show that the checkerboard copula has the largest Shannon entropy, which means that it carries the least information among all possible copulas for a given random vector. Furthermore, the checkerboard copula preserves the dependence information of the original random vector, leading to two applications in the context of diversification penalty and impact portfolios. The numerical and empirical results illustrate the benefits of using the checkerboard copula in the calculation of corisk measures.

Key words. orthant dependence, positive and negative association, Shannon entropy, CoVaR, simulation

MSC codes. 62H05, 91B30, 91G70

DOI. 10.1137/24M1656645

1. Introduction. The copula theory has been actively studied over the past few decades with many applications in statistics, finance, engineering, and the natural sciences; for an introduction, see the monographs of Nelsen (2006) and Joe (2015).

It is well known through Sklar's theorem (Nelsen (2006, Theorem 2.10.9)) that the copula of a random vector is unique if and only if it has continuous marginal distributions. However, when the marginals are noncontinuous, the uniqueness of the copula no longer holds. Discrete marginals are common in empirical studies, as the collected data are often discrete. Several works, including Marshall (1996), Carley (2002), Perrone, Solus, and Uhler (2019), and Geenens (2020), discuss the dependence structure of discrete data through copulas. Genest and Nešlehová (2007) discuss difficulties in identifying copulas for discrete distributions.

^{*}Received by the editors April 23, 2024; accepted for publication (in revised form) January 30, 2025; published electronically April 25, 2025.

https://doi.org/10.1137/24M1656645

Funding: The second author acknowledges financial support from the Natural Sciences and Engineering Research Council of Canada (RGPIN-2024-03728, CRC-2022-00141). The third author acknowledges research support from the National Key R&D Program of China (2022YFA1007900) and the National Natural Science Foundation of China (12271013, 72342004).

[†]Department of Statistics and Actuarial Science, University of Waterloo, Waterloo N2L 3G1, ON, Canada (I89lin@ uwaterloo.ca, wang@uwaterloo.ca).

[‡]Corresponding author. School of Mathematical Sciences, Center for Statistical Science, and National Engineering Laboratory for Big Data Analysis and Applications, Peking University, Beijing 100871, People's Republic of China (zhangruixun@pku.edu.cn).

[§]School of Mathematical Sciences, Peking University, Beijing 100871, People's Republic of China, and MIT Sloan School of Management and Laboratory for Financial Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139 USA (zhaochaoyi@pku.edu.cn).

The purpose of this paper is to understand whether it is possible to identify a canonical copula for a random vector in some sense if it does not have continuous marginal distributions.

To answer this question, we seek inspiration from four applications. Let $\mathbf{X} = (X_1, \ldots, X_d)$ be a *d*-dimensional random vector with $d \ge 2$, which may have nonunique copulas. Denote by $C_{\mathbf{X}}$ the set of all copulas of \mathbf{X} . For a random variable X, its probability integral transform U is a uniform random variable on [0,1] satisfying $F^{-1}(U) = X$ a.s., where F is the distribution function of X and F^{-1} is the quantile function of X. Let (U_1,\ldots,U_d) be any vector of probability integral transforms of X_1,\ldots,X_d with a joint distribution C; certainly, C is a copula of \mathbf{X} . All random variables live in an atomless probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

- 1. Simulating from the copula of X. One of the most popular applications of copulas in finance is to model default correlation, as famously done by Li (2000); see McNeil, Frey, and Embrechts (2015) for discussions. In such applications, one needs to simulate from the copula of X, where X may have noncontinuous marginal distributions (e.g., losses from default events). Assume that we can simulate X and that we also have knowledge of all marginal distributions of X. How can we find a reasonable copula $C \in C_X$ to simulate from, that is, determined only by X but not by any particular modeling choices (such as the Gaussian copula)?
- 2. Stressing the distribution of X. In sensitivity analysis and risk management, it is often necessary to stress, or distort, the distribution of X to obtain poststress distributions. In the stressing mechanisms studied by Millossovich, Tsanakas, and Wang (2024), one needs to find a stressed probability measure Q_1 by using $dQ_1/d\mathbb{P} = g(U_1)$ for a nonnegative increasing function g with $\int_0^1 g(u)du = 1$, such as g(u) = 2u. The simple interpretation of Q_1 is to gradually increase the weight of realizations $\omega \in \Omega$ at which X_1 is large. Similarly, one can simultaneously stress all components of X by considering a measure Q such that $dQ/d\mathbb{P} = (1/d) \sum_{i=1}^d g_i(U_i)$ or $dQ/d\mathbb{P} = c \prod_{i=1}^d g_i(U_i)$ with a normalizing constant c > 0 (c = 1 if U_1, \ldots, U_d are independent), where g_i are nonnegative increasing functions with $\int_0^1 g_i(u)du = 1$. If we are only interested in the poststress distribution $\hat{F}_1^{Q_1}$ of X_1 under Q_1 , then the choice of the copula $C \in C_X$ is irrelevant. However, the choice of the copula $C \in \mathcal{C}_X$ matters for the distribution \hat{F}_i^Q of X_i under Q as well as for the distribution $\hat{F}_i^{Q_1}$ of X_i under Q_1 .
- 3. Computing a corisk measure. Corisk measures (e.g., Adrian and Brunnermeier (2016)) are calculated for the conditional distribution of a random variable X_2 given some event related to X_1 . A classic example is the marginal expected shortfall (marginal ES) at level $p \in (0,1)$, which is defined as, assuming that X_1 is continuously distributed:

(1)
$$\rho(X_2|X_1) := \mathbb{E}[X_2|X_1 > F_1^{-1}(p)] = \mathbb{E}[X_2|U_1 > p].$$

Generally, ρ is the mean of X_2 given a (not necessarily unique) *p*-tail event of X_1 in the sense of Wang and Zitikis (2021). This risk measure ρ does not depend on the choice of $C \in \mathcal{C}_{\mathbf{X}}$ if X_1 is continuously distributed (*p*-tail event is unique a.s.); however, it may depend on $C \in \mathcal{C}_{\mathbf{X}}$ if X_1 has some points of mass. Other corisk measures, such as CoVaR (Adrian and Brunnermeier, 2016), also face the same issue.

4. Maintaining dependence measures. Kendall's τ is a dependence measure defined based on concordance. For a bivariate random vector (X, Y), its Kendall's τ is defined as

$$\tau(X,Y) = \mathbb{P}\left((X_1 - X_2)(Y_1 - Y_2) > 0\right) - \mathbb{P}\left((X_1 - X_2)(Y_1 - Y_2) < 0\right),$$

where (X_1, Y_1) and (X_2, Y_2) are two independent copies of (X, Y). If (X, Y) has continuous marginals, then we have further

(2)
$$\tau(X,Y) = 4 \int_{[0,1]^2} C(\mathbf{u}) dC(\mathbf{u}) - 1,$$

where C is the copula for (X, Y). If (X, Y) has noncontinuous marginals, then (2) does not hold for every $C \in \mathcal{C}_{(X,Y)}$. It is natural to ask which copula $C \in \mathcal{C}_{(X,Y)}$ conveys the property in the case of continuous marginals. We can also consider similar applications for other concordance-based dependence measures, such as Spearman's ρ .

All of the above contexts point to the question of choosing a good copula $C \in \mathcal{C}_{\mathbf{X}}$. Nešlehová (2004) discusses similar applications for the choice of copulas. The main idea of Nešlehová (2004) is to extend the subcopula, which is the part of the copula that is uniquely determined by joint distribution, to capture the dependence of the original random vector analogous to the case with continuous marginals. In this paper, we address this problem from the view of probability integral transformation. We first offer a new characterization of all copulas of a given random vector by constructing all probability integral transformations in section 2 in Theorem 1. In section 3, we give some intuitive and heuristic arguments for the questions above, leading to the proposal of using the checkerboard copula, that is, the unique copula of **X** that is as uniform as possible in regions where the copulas of **X** are not uniquely determined, formally defined in Definition 1. The checkerboard copula is the same as the standard extension proposed by Nešlehová (2004), which has been shown to preserve quadrant dependence, tail dependence, and weak convergence results of the joint distribution. Although the arguments in section 3 are heuristic, the use of the checkerboard copula indeed has a theoretical justification, which we present in section 4. The checkerboard copula has the maximum Shannon entropy among all possible copulas of \mathbf{X} , as shown in Theorem 2. In section 5, we show in Theorem 3 that the checkerboard copula preserves various dependence concepts that are satisfied by **X**. This result is intuitive, but the proof requires serious technical analysis. We discuss two applications of our results in diversification penalty and induced order statistics in section 6. Section 7 uses numerical and empirical experiments to demonstrate that the checkerboard copula is a convenient and natural choice that can produce reliable results. Section 8 concludes the paper.

2. Copulas for a discrete random vector. Let $d \ge 2$ be an integer, and let $[d] = \{1, \ldots, d\}$. All inequalities are interpreted componentwise when applied to vectors. All random variables live in an atomless probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $\mathbf{X} = (X_1, \ldots, X_d)$ be a *d*-dimensional random vector, F_1, \ldots, F_d be the marginal distributions of \mathbf{X} , and $\operatorname{Ran}(F_i)$ be the range of F_i for $i \in [d]$. By Sklar's theorem, the copula of \mathbf{X} is uniquely determined on $\operatorname{Ran}(F_1) \times \cdots \times$ $\operatorname{Ran}(F_d)$ but undetermined in other regions. Therefore, when the marginal distribution F_i is

not continuous for some $i \in [d]$, the copula of X may not be unique. In this section, we give a concrete representation for any copulas of X.

We start with the observation that if a random variable X is continuously distributed. then the random variable

$$U_X := F_X(X)$$

will be uniformly distributed over [0,1], where F_X is the cumulative distribution function of X. More generally, regardless of whether X is continuously distributed, we can define its probability integral transform

(3)
$$U_X := F_X(X-) + V_X(F_X(X) - F_X(X-)),$$

where $F_X(x-) = \lim_{y \uparrow x} F_X(x) = \mathbb{P}(X < x)$ for $x \in \mathbb{R}$ and $V_X \sim U[0,1]$ is independent of X, assumed to exist.¹ The probability integral transform U_X satisfies $U_X \sim U[0,1]$ and $F_X^{-1}(U_X) = X$ a.s. (see, e.g., Rüschendorf (2013, Proposition 1.3)). Therefore, the probability integral transform (3) converts any random variable X to a U[0,1] distributed random variable U_X using V_X .

We extend this idea to the case of a random vector **X**. Let $\mathbf{V} = (V_1, \ldots, V_d)$ be a random vector with U[0,1] marginals such that V_i is independent of X_i for each $i \in [d]$. Denote the set of such V by $\mathcal{V}_{\mathbf{X}}$. Similar to (3), let us define the probability integral transform for $\mathbf{X} = (X_1, \ldots, X_d):$

(4)
$$U_i := F_i(X_i) + V_i(F_i(X_i) - F_i(X_i)), \quad i \in [d].$$

It immediately follows that $U_i \sim U[0,1]$ and $F_i^{-1}(U_i) = X_i$ a.s. Therefore, $\mathbf{U} = (U_1, \ldots, U_d)$ is a random vector with uniform marginals. This transformation comes from randomized hypothesis tests (Ferguson (1967, section 5.3)) and has been applied in various contexts; see, e.g., Moore and Spruill (1975), Nešlehová (2007), Rüschendorf (1981, 2009, 2013), and Faugeras (2017).

Let $C_{\mathbf{X}}^{\mathbf{V}}$ be the copula of **U**. Because $F_i^{-1}(U_i) = X_i$ a.s. for each $i \in [d]$, we have

$$C_{\mathbf{X}}^{\mathbf{V}}(F_1(x_1),\ldots,F_d(x_d)) = \mathbb{P}(U_1 \leqslant F_1(x_1),\ldots,U_d \leqslant F_d(x_d)) = \mathbb{P}(X_1 \leqslant x_1,\ldots,X_d \leqslant x_d)$$

for any $(x_1, \ldots, x_d) \in \mathbb{R}^d$. Hence, $C_{\mathbf{X}}^{\mathbf{V}}$ is a copula of \mathbf{X} . According to (4), the copula $C_{\mathbf{X}}^{\mathbf{V}}$ is determined by the joint distribution of (\mathbf{X}, \mathbf{V}) . In particular, the copula $C_{\mathbf{X}}^{\mathbf{V}}$ does not depend on the choice of V_i for i such that X_i is continuously distributed because, for these i, U_i in (4) is a.s. equal to $F_i(X_i)$, while for i such that X_i is discrete, V_i does have an impact on the copula $C_{\mathbf{X}}^{\mathbf{V}}$.

In general, the choice of $\mathbf{V} \in \mathcal{V}_{\mathbf{X}}$ for constructing the copula $C_{\mathbf{X}}^{\mathbf{V}}$ may not be unique. This is because $\mathcal{V}_{\mathbf{X}}$ allows two types of dependence that might be present in the construction of V: First, the components of V may be mutually dependent. Second, V_i may depend on X_i

¹This assumption is safe, as we are interested in distributional properties, and we can extend the probability space to include such independent V_X , if necessary.

for $i \neq j$. Naturally, a different choice of $\mathbf{V} \in \mathcal{V}_{\mathbf{X}}$ often leads to a different copula $C_{\mathbf{X}}^{\mathbf{V}}$; see the following example.

Example 1. Assume that d = 2, X_1 is a constant and that X_2 is continuously distributed. It is well known that any copula is a copula of **X** in this case. For instance, by choosing V_1 to be independent of X_2 , $C_{\mathbf{X}}^{\mathbf{V}}$ is the independence copula, and by choosing $V_1 = F_2(X_2)$, $C_{\mathbf{X}}^{\mathbf{V}}$ is the comonotonic copula.

The following result says that all copulas of X can be realized by some $C_{\mathbf{X}}^{\mathbf{V}}$. Hence, (4) gives a stochastic representation for any copula of \mathbf{X} . The representation is quite intuitive, but we did not find it in the literature, so we provide a self-contained proof.

Theorem 1. Let \mathbf{X} be a random vector such that there exists a continuously distributed random variable independent of \mathbf{X} . A copula C is a copula of \mathbf{X} if and only if $C = C_{\mathbf{X}}^{\mathbf{V}}$ for some $\mathbf{V} \in \mathcal{V}_{\mathbf{X}}$.

Proof. We have seen that $C_{\mathbf{X}}^{\mathbf{V}}$ is a copula of \mathbf{X} . It suffices to show the "only if" statement. Let C be a copula of \mathbf{X} , take $\mathbf{U}' = (U'_1, \ldots, U'_d) \sim C$, and write $\mathbf{X}' = (F_1^{-1}(U'_1), \ldots, F_d^{-1}(U'_d))$. Because C is a copula of \mathbf{X} , for $\mathbf{x} = (x_1, \ldots, x_d) \in \mathbb{R}^d$, we have

$$\mathbb{P}(\mathbf{X} \leq \mathbf{x}) = C\left(F_1(x_1), \dots, F_d(x_d)\right) = \mathbb{P}\left(U_1' \leq F_1(x_1), \dots, U_d' \leq F_d(x_d)\right)$$
$$= \mathbb{P}\left(F_1^{-1}(U_1') \leq x_1, \dots, F_d^{-1}(U_d') \leq x_d\right) = \mathbb{P}(\mathbf{X}' \leq \mathbf{x}).$$

Hence, $\mathbf{X} \stackrel{d}{=} \mathbf{X}'$. Take $\mathbf{U}^* = (U_1^*, \dots, U_d^*)$ such that $(\mathbf{X}, \mathbf{U}^*) \stackrel{d}{=} (\mathbf{X}', \mathbf{U}')$, and we then have $\mathbf{X} = (F_1^{-1}(U_1^*), \dots, F_d^{-1}(U_d^*))$ a.s. Furthermore, take $V' \sim U[0, 1]$, which is independent of $(\mathbf{X}, \mathbf{U}^*)$. The existence of \mathbf{U}^* and V' is guaranteed by the assumption of the existence of a continuously distributed random variable independent of \mathbf{X} . For $i \in [d]$, let $\mathbf{V} = (V_1, \dots, V_d)$ be given by

$$V_{i} = \frac{U_{i}^{*} - F_{i}(X_{i})}{F_{i}(X_{i}) - F_{i}(X_{i})} \mathbb{1}_{\{F_{i}(X_{i}) > F_{i}(X_{i})\}} + V' \mathbb{1}_{\{F_{i}(X_{i}) = F_{i}(X_{i})\}}.$$

Fix $i \in [d]$ below. Let D_i be the set of discontinuity points of F_i . Note that for $x \in D_i$, we have

$$\mathbb{P}(U_i^* \in [F_i(x-), F_i(x)] | X_i = x) = 1 \text{ and } \mathbb{P}(U_i^* \in [F_i(x-), F_i(x)] | X_i \neq x) = 0.$$

Because U_i^* is uniformly distributed over [0,1], U_i^* is uniform on $[F_i(x-), F_i(x)]$ conditional on $X_i = x \in D_i$. Thus,

$$\mathbb{P}(U_i^* \leqslant u | X_i = x) = \frac{u - F_i(x-)}{F_i(x) - F_i(x-)}, \quad u \in [F_i^{-1}(x-), F_i^{-1}(x)].$$

Therefore, for $u \in [0, 1]$,

$$\mathbb{P}(V_i \leqslant u | X_i) = \mathbb{P}\left(\frac{U_i^* - F_i(X_i)}{F_i(X_i) - F_i(X_i)} \leqslant u | X_i\right) \mathbb{1}_{\{X_i \in D_i\}} + \mathbb{P}(V' \leqslant u) \mathbb{1}_{\{X_i \notin D_i\}}$$

= $\mathbb{P}\left(U_i^* \leqslant u(F_i(X_i) - F_i(X_i)) + F_i(X_i) \mathbb{1}_{\{X_i \in D_i\}} + u \mathbb{1}_{\{X_i \notin D_i\}}$
= $u \mathbb{1}_{\{X_i \in D_i\}} + u \mathbb{1}_{\{X_i \notin D_i\}} = u.$

Hence, V_i follows U[0,1] and is independent of X_i . Note that, by the construction, U_i^* , V_i , and X_i satisfy $U_i^* = F_i(X_i-) + V_i(F_i(X_i) - F_i(X_i-))$ a.s., and hence $\mathbf{U}^* \sim C_{\mathbf{X}}^{\mathbf{V}}$. This shows that $C = C_{\mathbf{X}}^{\mathbf{V}}$.

Theorem 1 implies that $C_{\mathbf{X}} = \{C_{\mathbf{X}}^{\mathbf{V}} : \mathbf{V} \in \mathcal{V}_{\mathbf{X}}\}$, providing a characterization of copulas in a stochastic form. Note that $C_{\mathbf{X}}$ is a singleton if and only if all marginal distributions of \mathbf{X} , F_1, \ldots, F_d , are continuous functions. The "if" direction of Theorem 1 in the case d = 2 is shown by Nešlehová (2007, Proposition 4). The characterization of copulas in an analytical form is provided by de Amo et al. (2017).

3. Motivating arguments for the checkerboard copula. Theorem 1 gives the entire class of copulas for **X**. We now consider which $\mathbf{V} \in \mathcal{V}_{\mathbf{X}}$ can answer the four motivating questions in section 1, which all point to the same unique choice of $\mathbf{V} \in \mathcal{V}_{\mathbf{X}}$.

- 1. Simulating from the copula of X. A natural approach to simulating from the copula of X with some atoms in the marginal distributions is by first simulating a pair of (\mathbf{X}, \mathbf{V}) and then applying the probability integral transform using (4). Theorem 1 shows that all copulas of X can be simulated this way. For this purpose, the simplest and most natural choice of \mathbf{V} is $\mathbf{V} \sim \mathrm{U}([0,1]^d)$, which is independent of X. In fact, we could not think of an argument against the use of this particular V in the context of simulation.
- 2. Stressing the distribution of X. To understand how the choice of V affects the stressed distribution of X_2 , we look at the simple example in Example 1 with g(u) = 2u. Choosing V_1 independent of X_2 would lead to $\hat{F}_2^{Q_1} = F_2$, whereas choosing $V_1 = F_2(X_2)$ would lead to $\hat{F}_2^{Q_1} = (F_2)^2$. Because we are interested in the effect of stressing X_1 on X_2 and because X_1 is a constant in this example, it is natural to choose a V_1 that affects the distribution of X_2 minimally, which is achieved when V_1 is independent of X_2 . Translating this argument into the general *d*-dimensional setting suggests choosing $\mathbf{V} \sim \mathbf{U}([0,1]^d)$ independent of \mathbf{X} .
- 3. Computing a corisk measure. To understand how the choice of V affects the value of the corisk measure, we again look at Example 1. We have $\rho(X_2|X_1) = \mathbb{E}[X_2]$ if V_1 is independent of X_2 and $\rho(X_2|X_1) = \mathrm{ES}_p(X_2)$ if $V_1 = F_2(X_2)$, where $\mathrm{ES}_p(X_2) = \mathbb{E}[X_2|U_2 > p]$ is the ES of X_2 at level p. The interpretation of ρ as the mean of X_2 on a tail event of X_1 suggests that it is natural to choose V_1 independent of X_2 because X_1 is a constant and its tail event should not affect X_2 .
- 4. Maintaining dependence measures. It has been shown in Denuit and Lambert (2005), Nešlehová (2007), and Genest and Nešlehová (2007) that (2) holds for the copula introduced by $\mathbf{V} \sim \mathrm{U}([0,1]^2)$. This property leads to extensions of dependence measures in the multivariate case based on this copula; see, e.g., Mesfioui and Quessy (2010) and Genest, Nešlehová, and Rémillard (2013).

In all the considerations above, $\mathbf{V} \sim \mathrm{U}([0,1]^d)$ independent of \mathbf{X} appears to be a good choice. Let us denote this by $\mathbf{V}_{\mathbf{X}}^{\perp}$ and the corresponding copula by $C_{\mathbf{X}}^{\perp}$, where \perp reflects that independence is used twice to construct \mathbf{V} (within components of \mathbf{V} and between \mathbf{V} and \mathbf{X}). From the four motivating examples above, the choice of the particular copula $C_{\mathbf{X}}^{\perp}$ is natural and has several unique features. This choice has been known as the checkerboard copula.

Definition 1. The copula $C_{\mathbf{X}}^{\perp}$ is called the checkerboard copula of \mathbf{X} .

The copula $C_{\mathbf{X}}^{\perp}$ is also called the multilinear extension copula of \mathbf{X} ; see Genest, Nešlehová, and Rémillard (2017) for its properties, its empirical process, and a history. One notable property is that X_1, \ldots, X_d are independent if and only if $C_{\mathbf{X}}^{\perp}$ is the independence copula.

The rest of the paper focuses on the properties and applications of the checkerboard copula.

4. Entropy maximization. Given the natural choice of $C_{\mathbf{X}}^{\perp}$ in the applications in section 3, it should have some unique properties within the class $C_{\mathbf{X}}$. The applications seem to suggest that $C_{\mathbf{X}}^{\perp}$ relies less on external information compared to other choices of **V**. Such consideration is typically studied via entropy. Indeed, as argued by Jaynes (1957), the maximum-entropy distribution should be the only unbiased choice given the available information. If a copula C has a density function c, then its Shannon (differential) entropy is defined as

$$H(C) = -\int_{[0,1]^d} c(\mathbf{u}) \log c(\mathbf{u}) \mathrm{d}\mathbf{u}.$$

One problem with the above formulation is that a copula C often does not have a density. We set $H(C) = -\infty$ if C does not have a density, which is intuitive and can be seen as a limiting case; see Koliander et al. (2016) for a discussion on the definition of entropy for singular distributions.

Remark 1. By definition, $H(C) = -D_{\mathrm{KL}}(P_C || P_L)$, where $D_{\mathrm{KL}}(P_C || P_L)$ is the Kullback– Leibler (KL) divergence between the probability measure P_C with distribution function Cand the Lebesgue measure P_L on $[0,1]^d$. Since the KL divergence quantifies the similarity between P_C and P_L via the likelihood ratio dP_C/dP_L , being singular is the extreme form of nonsimilarity in terms of likelihood ratio. Therefore, it is natural to set $D_{\mathrm{KL}}(P_C || P_L)$ as ∞ whenever P_C is not absolutely continuous with respect to P_L . This is a standard approach in the literature on KL divergence and differential entropy; see, e.g., Csiszár (1975) and Cover and Thomas (1991). Hence, to keep the same intuition and consistency with the KL divergence, we set $H(C) = -\infty$ when C does not have density.

However, even the checkerboard copula $C_{\mathbf{X}}^{\perp}$ may not have a density if the distribution of **X** has some singular continuous part. This issue may be solved by considering other measures of information, but for now, let us stick to the Shannon entropy, which is the most popular notion in information theory. We would like to compare $H(C_{\mathbf{X}}^{\perp})$ with H(C) for $C \in \mathcal{C}_{\mathbf{X}}$ or, equivalently, $H(C_{\mathbf{X}}^{\mathbf{V}})$ for other choices of $\mathbf{V} \in \mathcal{V}_{\mathbf{X}}$. The main result of this section is to show that $H(C_{\mathbf{X}}^{\perp})$ has the largest entropy among all other choices.

Theorem 2. For $C \in \mathcal{C}_{\mathbf{X}}$, we have $H(C_{\mathbf{X}}^{\perp}) \geq H(C)$.

The proof of Theorem 2 essentially boils down to showing the following lemma, which states that the density of the checkerboard copula can be expressed as the conditional expectation for the density of other possible copulas in $C_{\mathbf{X}}$. From this lemma and Jensen's inequality, Theorem 2 follows.

Lemma 1. For $C \in \mathcal{C}_{\mathbf{X}}$, if the density c of C exists, then the density c^{\perp} of $C_{\mathbf{X}}^{\perp}$ exists. Moreover, $c^{\perp}(\mathbf{U}) = \mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}]$, where $\mathbf{U} = (U_1, \ldots, U_d) \sim \mathrm{U}([0,1]^d)$, $\hat{\mathbf{X}} = (F_1^{-1}(U_1), \ldots, F_d^{-1}(U_d))$, and F_1, \ldots, F_d are the marginals of \mathbf{X} . *Proof.* Since $\mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}]$ is $\sigma(\hat{\mathbf{X}})$ -measurable, there exists a function $f: \mathbb{R}^d \to [0,1]$ such that $f(\hat{\mathbf{X}}) = \mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}]$ (in the almost sure sense). Let c^{\perp} be a function $[0,1]^d \to [0,1]$ defined as $c^{\perp}(\mathbf{u}) = f\left(F_1^{-1}(u_1), \ldots, F_d^{-1}(u_d)\right)$ for any $\mathbf{u} = (u_1, \ldots, u_d) \in [0,1]^d$. We claim that c^{\perp} is the density of $C_{\mathbf{X}}^{\perp}$. This claim implies that c^{\perp} exists and that $c^{\perp}(\mathbf{U}) = \mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}]$.

density of $C_{\mathbf{X}}^{\perp}$. This claim implies that c^{\perp} exists and that $c^{\perp}(\mathbf{U}) = \mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}]$. To prove this claim, let $\mathbf{U}^c \sim C$, $\mathbf{U}^{\perp} \sim C_{\mathbf{X}}^{\perp}$, and $R = \mathbf{X}_{i=1}^d \operatorname{Ran}(F_i)$. We first show that $\int_A c^{\perp}(\mathbf{u}) d\mathbf{u} = \mathbb{P}(\mathbf{U}^{\perp} \in A)$ for the following two types of the set A.

(i) $A = \bigotimes_{i=1}^{d} [0, a_i]$ with $\mathbf{a} = (a_1, \dots, a_d) \in R$. We have $\mathbb{1}_{\{\mathbf{U} \leq \mathbf{a}\}} = \mathbb{1}_{\{\hat{\mathbf{X}} \leq (F_1^{-1}(a_1), \dots, F_d^{-1}(a_d))\}}$. Therefore,

$$\begin{split} \int_{A} c^{\perp}(\mathbf{u}) \mathrm{d}\mathbf{u} &= \int_{\times_{i=1}^{d}[0,a_{i}]} f\left(F_{1}^{-1}(u_{1}), \dots, F_{d}^{-1}(u_{d})\right) \mathrm{d}u_{1} \cdots \mathrm{d}u_{d} \\ &= \mathbb{E}\left[f(\hat{\mathbf{X}})\mathbb{1}_{\{\mathbf{U} \leq \mathbf{a}\}}\right] \\ &= \mathbb{E}\left[\mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}]\mathbb{1}_{\{\hat{\mathbf{X}} \leq \left(F_{1}^{-1}(a_{1}), \dots, F_{d}^{-1}(a_{d})\right)\}}\right] \\ &= \mathbb{E}\left[c(\mathbf{U})\mathbb{1}_{\{\mathbf{U} \leq \mathbf{a}\}}\right] = \mathbb{P}(\mathbf{U}^{c} \leq \mathbf{a}) = \mathbb{P}(\mathbf{U}^{\perp} \leq \mathbf{a}), \end{split}$$

where the last equality holds because

$$\mathbb{P}(\mathbf{U}^c \leq \mathbf{a}) = \mathbb{P}\left(\mathbf{X} \leq \left(F_1^{-1}(a_1), \dots, F_d^{-1}(a_d)\right)\right) = \mathbb{P}(\mathbf{U}^{\perp} \leq \mathbf{a}).$$

This further implies that $\int_A c^{\perp}(\mathbf{u}) d\mathbf{u} = \mathbb{P}(\mathbf{U}^{\perp} \in A)$ for any $A = \bigotimes_{i=1}^d A_i$ such that $A_i \in \{[0, a_i] : a_i \in \operatorname{Ran}(F_i)\} \cup \{(F_i(x_i-), F_i(x_i)] : x_i \text{ is a discontinuity point of } F_i\}$ for $i \in [d]$.

(ii) Let $A = (\bigotimes_{i=1}^{k} [0, a_i]) \times (\bigotimes_{j=k+1}^{d} (s_j, t_j])$ with $k \in \{0, 1, \dots, d\}$ such that $a_i \in \operatorname{Ran}(F_i)$ for $i \in [k]$ and $(s_j, t_j] \cap \operatorname{Ran}(F_j) = \emptyset$ for $j \in [d] \setminus [k]$. For $j \in [d] \setminus [k]$, denote by $x_j = F_j^{-1}(s_j)$, and thus $(s_j, t_j] \subseteq (F_j(x_j -), F_j(x_j))$. By the definition of c^{\perp} , for fixed $u_i \in [0, a_i]$ and $i \in [k]$, $c^{\perp}(u_1, \dots, u_k, v_{k+1}, \dots, v_d)$ is a constant for all $(v_{k+1}, \dots, v_d) \in$ $\bigotimes_{j=k+1}^{d} (F_j(x_j -), F_j(x_j))$. Therefore, let $B = (\bigotimes_{i=1}^{k} [0, a_i]) \times (\bigotimes_{j=k+1}^{d} (F_j(x_j -), F_j(x_j)))$. We have

$$\int_{A} c^{\perp}(\mathbf{u}) \mathrm{d}\mathbf{u} = \left(\prod_{j=k+1}^{d} \frac{t_j - s_j}{F_j(x_j) - F_j(x_j)}\right) \int_{B} c^{\perp}(\mathbf{u}) \mathrm{d}\mathbf{u}$$

Let $\mathbf{V} = (V_1, ..., V_d) \sim U([0, 1]^d)$ be independent of \mathbf{X} , and for $j \in [d] \setminus [k]$, denote by $s'_j = (s_j - F_j(x_j -))/(F_j(x_j) - F_j(x_j -))$ and $t'_j = (t_j - F_j(x_j -))/(F_j(x_j) - F_j(x_j -))$. Hence,

$$\prod_{j=k+1}^{d} \frac{t_j - s_j}{F_j(x_j) - F_j(x_j)} = \mathbb{P}\left(V_j \in (s'_j, t'_j] \text{ for all } j \in [d] \setminus [k]\right).$$

In addition, by (i), we can get

$$\int_{B} c^{\perp}(\mathbf{u}) \mathrm{d}\mathbf{u} = \mathbb{P}(\mathbf{U}^{\perp} \in B) = \mathbb{P}\left(X_{i} \leqslant F^{-1}(a_{i}), X_{j} = x_{j} \text{ for all } i \in [k], j \in [d] \setminus [k]\right).$$

Therefore,

$$\begin{split} \int_{A} c^{\perp}(\mathbf{u}) \mathrm{d}\mathbf{u} &= \mathbb{P}\left(\bigcap_{j \in [d] \setminus [k]} \{V_{j} \in (s'_{j}, t'_{j}]\}\right) \mathbb{P}\left(\bigcap_{i \in [k], j \in [d] \setminus [k]} \{X_{i} \leqslant F^{-1}(a_{i}), \ X_{j} = x_{j}\}\right) \\ &= \mathbb{P}\left(\bigcap_{i \in [k], j \in [d] \setminus [k]} \{X_{i} \leqslant F^{-1}(a_{i}), \ X_{j} = x_{j}, \ V_{i} \in [0, 1], \ V_{j} \in (s'_{j}, t'_{j}]\}\right) \\ &= \mathbb{P}\left(\mathbf{U}^{\perp} \in \left(\bigotimes_{i=1}^{k} [0, a_{i}]\right) \times \left(\bigotimes_{j=k+1}^{d} (s_{j}, t_{j}]\right)\right) = \mathbb{P}(\mathbf{U}^{\perp} \in A). \end{split}$$

By the same argument, we have $\int_A c^{\perp}(\mathbf{u}) d\mathbf{u} = \mathbb{P}(\mathbf{U}^{\perp} \in A)$ for any $A = \mathbf{X}_{i=1}^d A_i$ such that $A_i \in \{[0, a_i] : a_i \in \operatorname{Ran}(F_i)\} \cup \{(s_i, t_i] : (s_i, t_i] \cap \operatorname{Ran}(F_i) = \emptyset\}$ for $i \in [d]$.

For any $\mathbf{a} = (a_1, \ldots, a_d) \in [0, 1]^d$, the region $\times_{i=1}^d [0, a_i]$ can always be represented by an at most countable disjoint union of regions studied in (i) and (ii). Hence, we can obtain

$$\int_{\times_{i=1}^{d}[0,a_i]} c^{\perp}(\mathbf{u}) \mathrm{d}\mathbf{u} = \mathbb{P}(\mathbf{U}^{\perp} \leqslant \mathbf{a}).$$

This proves our claim that c^{\perp} is the density of $C_{\mathbf{x}}^{\perp}$.

Proof of Theorem 2. If $H(C) = -\infty$, then there is nothing to show. Hence, it suffices to consider the case that C has a density, which we denote by c. By Lemma 1, we have $c^{\perp}(\mathbf{U}) = \mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}]$, where c^{\perp} is the density of $C_{\mathbf{X}}^{\perp}$, $\mathbf{U} = (U_1, \ldots, U_d) \sim \mathbf{U}([0, 1]^d)$, and $\hat{\mathbf{X}} = (F_1^{-1}(U_1), \ldots, F_d^{-1}(U_d))$ with F_1, \ldots, F_d as the marginals of \mathbf{X} . Define a function $g(x) = x \log x$ for $x \in (0, \infty)$. It is clear that g is convex. By the fact that $\mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}] = c^{\perp}(\mathbf{U})$ and Jensen's inequality, we have

$$H(C_{\mathbf{X}}^{\perp}) = -\mathbb{E}[g(c^{\perp}(\mathbf{U}))] = -\mathbb{E}[g(\mathbb{E}[c(\mathbf{U})|\hat{\mathbf{X}}])] \ge -\mathbb{E}[\mathbb{E}[g(c(\mathbf{U}))|\hat{\mathbf{X}}]] = -\mathbb{E}[g(c(\mathbf{U}))] = H(C).$$

Thus, $H(C_{\mathbf{X}}^{\perp}) \geq H(C)$ for all $C \in \mathcal{C}_{\mathbf{X}}$.

Theorem 2 demonstrates that the entropy of $C_{\mathbf{X}}$ cannot be greater than $C_{\mathbf{X}}^{\perp}$. This result is related to those of Piantadosi, Howlett, and Borwein (2012) and Kuzmenko, Salam, and Uryasev (2020), and the difference is that, in Theorem 2, we fix a joint (possibly discrete) distribution and seek to find the copula consistent with this distribution that maximizes the entropy, which is the checkerboard copula. In contrast, Piantadosi, Howlett, and Borwein (2012) and Kuzmenko, Salam, and Uryasev (2020) do not fix a joint distribution. Instead, they search for a checkerboard copula that maximizes the entropy subject to matching either a correlation coefficient or the distribution of the sum of random variables. Therefore, the problems they address are different from our Theorem 2.

When $C_{\mathbf{X}}$ and $C_{\mathbf{X}}^{\perp}$ contain singular components, by definition, $H(C_{\mathbf{X}}) = H(C_{\mathbf{X}}^{\perp}) = -\infty$. In this case, the next proposition shows that the entropy for the absolutely continuous part of $C_{\mathbf{X}}^{\perp}$ is still greater than that for the absolutely continuous part of $C_{\mathbf{X}}$.

THE CHECKERBOARD COPULA AND DEPENDENCE CONCEPTS

Proposition 1. Assume that $C_{\mathbf{X}} \in \mathcal{C}_{\mathbf{X}}$ such that $C_{\mathbf{X}} = \lambda G_{\mathrm{A}} + (1-\lambda)G_{\mathrm{S}}$, where $\lambda \in [0,1]$, G_{A} is an absolutely continuous distribution function and G_{S} is a singular distribution function. Then there exists an absolutely continuous distribution function G_{A}^{\perp} and a distribution function G_{S}^{\perp} such that $C_{\mathbf{X}}^{\perp} = \lambda G_{\mathrm{A}}^{\perp} + (1-\lambda)G_{\mathrm{S}}^{\perp}$ and $H(G_{\mathrm{A}}^{\perp}) \ge H(G_{\mathrm{A}})$.

Proof. Let F_1, \ldots, F_d be the marginal distributions of **X**. For $x \in [0, 1]$, let $l_i(x) = \sup\{y : y \in \operatorname{Ran}(F_i), y \leq x\}$ and $u_i(x) = \inf\{y : y \in \operatorname{Ran}(F_i), y \geq x\}$. Define two distribution functions G_A^{\perp} and G_S^{\perp} , which are linear interpolations of G_A and G_S from $\times_{i=1}^d \operatorname{Ran}(F_i)$ to $[0, 1]^d$: For $\mathbf{x} = (x_1, \ldots, x_d) \in [0, 1]^d$,

$$G_{\mathbf{A}}^{\perp}(\mathbf{x}) = \begin{cases} G_{\mathbf{A}}(\mathbf{x}), & \mathbf{x} \in \bigwedge_{i=1}^{d} \operatorname{Ran}(F_{i}), \\ \sum_{\substack{y_{i} \in \{l_{i}(x_{i}), u_{i}(x_{i})\} \\ i \in [d]}} \prod_{j=1}^{d} \beta_{j}(x_{j}, y_{j}) G_{\mathbf{A}}(y_{1}, \dots, y_{d}), & \mathbf{x} \in [0, 1]^{d} \setminus \bigotimes_{i=1}^{d} \operatorname{Ran}(F_{i}), \end{cases}$$

and

$$G_{\mathbf{S}}^{\perp}(\mathbf{x}) = \begin{cases} G_{\mathbf{S}}(\mathbf{x}), & \mathbf{x} \in \bigotimes_{i=1}^{d} \operatorname{Ran}(F_{i}), \\ \sum_{\substack{y_{i} \in \{l_{i}(x_{i}), u_{i}(x_{i})\} \\ i \in [d]}} \prod_{j=1}^{d} \beta_{j}(x_{j}, y_{j}) G_{\mathbf{S}}(y_{1}, \dots, y_{d}), & \mathbf{x} \in [0, 1]^{d} \setminus \bigotimes_{i=1}^{d} \operatorname{Ran}(F_{i}), \end{cases}$$

where

$$\beta_i(x,y) = \frac{u_i(x) - x}{u_i(x) - l_i(x)} \mathbb{1}_{\{y = l_i(x)\}} + \frac{x - l_i(x)}{u_i(x) - l_i(x)} \left(1 - \mathbb{1}_{\{y = l_i(x)\}}\right)$$

for $x, y \in [0,1]$ and $i \in [d]$ with the convention 0/0 = 1. Note that $\beta_i(x, y)$ is linear in x on each segment of $[0,1] \setminus \text{Ran}(F_i)$. It is clear that G_A^{\perp} is continuous.

Let g_A be the density of G_A and g be the derivative of G_A^{\perp} , respectively. For $\mathbf{x} \in \mathbf{x}_{i=1}^d \operatorname{Ran}(F_i)$, we have

$$g(\mathbf{x}) = \frac{\partial^d G_{\mathbf{A}}^{\perp}(\mathbf{x})}{\partial x_1 \cdots \partial x_d} = \frac{\partial^d G_{\mathbf{A}}(\mathbf{x})}{\partial x_1 \cdots \partial x_d} = g_{\mathbf{A}}(\mathbf{x}).$$

For $\mathbf{x} = (x_1, \dots, x_{d-1}, x_d)$ such that $x_i \in \operatorname{Ran}(F_i)$ for $i \in [d-1]$ and $x_d \notin \operatorname{Ran}(F_d)$, we have

$$G_{\mathcal{A}}^{\perp}(\mathbf{x}) = \frac{u_d(x_d) - x_d}{u_d(x_d) - l_d(x_d)} G_{\mathcal{A}}(x_1, \dots, x_{d-1}, l_d(x_d)) + \frac{x_d - l_d(x_d)}{u_d(x_d) - l_d(x_d)} G_{\mathcal{A}}(x_1, \dots, x_{d-1}, u_d(x_d)).$$

Hence,

$$g(\mathbf{x}) = \frac{\partial^d G_{\mathbf{A}}^{\perp}(\mathbf{x})}{\partial x_1 \cdots \partial x_d}$$

= $\frac{\partial^{d-1} [G_{\mathbf{A}}(x_1, \dots, x_{d-1}, u_d(x_d)) - G_{\mathbf{A}}(x_1, \dots, x_{d-1}, l_d(x_d))]}{\partial x_1 \cdots \partial x_{d-1}}$
= $\int_{l_d(x_d)}^{u_d(x_d)} g_{\mathbf{A}}(x_1, \dots, x_{d-1}, y) \mathrm{d}y.$

Let $k \in [d]$. Similarly, for $\mathbf{x} = (x_1, \dots, x_d)$ such that $x_i \in \operatorname{Ran}(F_i)$ for $i \in [k-1]$ (with $[0] = \emptyset$) and $x_i \notin \operatorname{Ran}(F_i)$ for $i \in [d] \setminus [k-1]$, we have

$$g(\mathbf{x}) = \int_{l_k(x_k)}^{u_k(x_k)} \dots \int_{l_d(x_d)}^{u_d(x_d)} g_{\mathbf{A}}(x_1, \dots, x_{k-1}, y_k, \dots, y_d) \mathrm{d}y_k \cdots \mathrm{d}y_d.$$

Note that $u_i(x_i) = F_i(F_i^{-1}(x_i))$ and that $l_i(x_i) = F_i(F_i^{-1}(x_i))$ for $i \in [d]$. From the discussion above, we can see that $g(\mathbf{x})$ is a constant in $\times_{i=1}^d [F_i(y_i-), F_i(y_i)]$ for all $\mathbf{y} = (y_1, \ldots, y_d) \in \mathbb{R}^d$. Let $\mathbf{U} = (U_1, \ldots, U_d) \sim \mathrm{U}([0, 1]^d)$, and let $\hat{\mathbf{X}} = (F_1^{-1}(U_1), \ldots, F_d^{-1}(U_d))$. Thus, $g(\mathbf{U}) = \mathbb{E}[g_{\mathbf{A}}(\mathbf{U})|\hat{\mathbf{X}}]$. Hence, g is Lebesgue integrable, and $G_{\mathbf{A}}^{\perp}$ is an absolutely continuous distribution function with density function $g_{\mathbf{A}}^{\perp} = g$.

Next, we show that $C_{\mathbf{X}}^{\perp} = \lambda G_{\mathbf{A}}^{\perp} + (1-\lambda)G_{\mathbf{S}}^{\perp}$ and $H(G_{\mathbf{A}}^{\perp}) \ge H(G_{\mathbf{A}})$. For $\mathbf{x} \in \mathbf{X}_{i=1}^{d} \operatorname{Ran}(F_{i})$,

$$\lambda G_{\mathcal{A}}^{\perp}(\mathbf{x}) + (1-\lambda)G_{\mathcal{S}}^{\perp}(\mathbf{x}) = \lambda G_{\mathcal{A}}(\mathbf{x}) + (1-\lambda)G_{\mathcal{S}}(\mathbf{x}) = C_{\mathbf{X}}(\mathbf{x}) = C_{\mathbf{X}}^{\perp}(\mathbf{x})$$

For $\mathbf{x} \in [0,1]^d \setminus \bigotimes_{i=1}^d \operatorname{Ran}(F_i)$,

$$\lambda G_{\mathcal{A}}^{\perp}(\mathbf{x}) + (1-\lambda)G_{\mathcal{S}}^{\perp}(\mathbf{x}) = \sum_{\substack{y_i \in \{l_i(x_i), u_i(x_i)\} \\ i \in [d]}} \prod_{j=1}^d \beta(x_j, y_j) \left(\lambda G_{\mathcal{A}}(y_1, \dots, y_d) + (1-\lambda)G_{\mathcal{S}}(y_1, \dots, y_d)\right)$$
$$= \sum_{\substack{y_i \in \{l_i(x_i), u_i(x_i)\} \\ i \in [d]}} \prod_{j=1}^d \beta(x_j, y_j) C_{\mathbf{X}}(y_1, \dots, y_d) = C_{\mathbf{X}}^{\perp}(\mathbf{x}).$$

Let $h(x) = -x \log x$ for $x \in (0, \infty)$. It is clear that h is a concave function. By Jensen's inequality, we have

$$H(G_{\mathbf{A}}^{\perp}) = \mathbb{E}[h(g_{\mathbf{A}}^{\perp}(\mathbf{U}))] = \mathbb{E}[h(\mathbb{E}[g_{\mathbf{A}}(\mathbf{U})|\hat{\mathbf{X}}])] \ge \mathbb{E}[\mathbb{E}[h(g_{\mathbf{A}}(\mathbf{U}))|\hat{\mathbf{X}}]] = \mathbb{E}[h(g_{\mathbf{A}}(\mathbf{U}))] = H(G_{\mathbf{A}}).$$

This completes the proof.

5. Checkerboard copula and dependence concepts. In this section, we study how the checkerboard copula preserves dependence concepts. This question is motivated by a problem raised in the context of diversification in Chen, Embrechts, and Wang (2024b), which we describe in section 6.1.

5.1. Dependence concepts. We first define several notions of positive dependence, introduced and studied by Lehmann (1966), Esary, Proschan, and Walkup (1967), and Benjamini and Yekutieli (2001), and the corresponding notions of negative dependence, introduced and studied by Lehmann (1966), Alam and Saxena (1981), Block, Savits, and Shaked (1982, 1985), Joag-Dev and Proschan (1983), and Chen, Embrechts, and Wang (2024a).

In what follows, for $i \in [d]$ and a *d*-dimensional random vector $\mathbf{X} = (X_1, \ldots, X_d)$, write $\mathbf{X}_{-i} = (X_1, \ldots, X_{i-1}, X_{i+1}, \ldots, X_d)$, and for $A, B \subseteq [d]$, write $\mathbf{X}_A = (X_k)_{k \in A}$ and $\mathbf{X}_B = (X_k)_{k \in B}$. A set $S \subseteq \mathbb{R}^d$ is *decreasing* if $\mathbf{x} \in S$ implies that $\mathbf{y} \in S$ for all $\mathbf{y} \leq \mathbf{x}$.

Copyright © by SIAM. Unauthorized reproduction of this article is prohibited.

Definition 2. A random vector \mathbf{X} is

 (i) (a) positively associated (PA) if for every pair of subsets A, B of [d] and any functions f and g both increasing or decreasing coordinatewise, provided that the covariance below exists,

$$\operatorname{Cov}(f(\mathbf{X}_A), g(\mathbf{X}_B)) \ge 0;$$

(b) negatively associated (NA) if for every pair of disjoint subsets A, B of [d] and any functions f and g both increasing or decreasing coordinatewise, provided that the covariance below exists,

$$\operatorname{Cov}(f(\mathbf{X}_A), g(\mathbf{X}_B)) \leq 0;$$

- (ii) (a) positively regression dependent (*PRD*) if for every $i \in [d]$, the random variable $\mathbb{E}[g(\mathbf{X}_{-i})|X_i]$ is an increasing function of X_i for any coordinatewise increasing function g such that the conditional expectation exists;
 - (b) negatively regression dependent (NRD) if for every $i \in [d]$, the random variable $\mathbb{E}[g(\mathbf{X}_{-i})|X_i]$ is a decreasing function of X_i for any coordinatewise increasing function g such that the conditional expectation exists;
- (iii) (a) weakly positively associated *(WPA)* if for any $i \in [d]$, decreasing set $S \subseteq \mathbb{R}^{d-1}$, and $x \in \mathbb{R}$ with $\mathbb{P}(X_i \leq x) > 0$,

$$\mathbb{P}(\mathbf{X}_{-i} \in S \mid X_i \leqslant x) \ge \mathbb{P}(\mathbf{X}_{-i} \in S);$$

(b) weakly negatively associated (WNA) if for any $i \in [d]$, decreasing set $S \subseteq \mathbb{R}^{d-1}$, and $x \in \mathbb{R}$ with $\mathbb{P}(X_i \leq x) > 0$,

$$\mathbb{P}(\mathbf{X}_{-i} \in S \mid X_i \leqslant x) \leqslant \mathbb{P}(\mathbf{X}_{-i} \in S);$$

- (iv) (a) positively orthant dependent *(POD)* if for all $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$, $\mathbb{P}(\mathbf{X} \leq \mathbf{x}) \geq \prod_{i=1}^d \mathbb{P}(X_i \leq x_i)$ and $\mathbb{P}(\mathbf{X} > \mathbf{x}) \geq \prod_{i=1}^d \mathbb{P}(X_i > x_i)$;
 - (b) negatively orthant dependent (NOD) if for all $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$, $\mathbb{P}(\mathbf{X} \leq \mathbf{x}) \leq \prod_{i=1}^d \mathbb{P}(X_i \leq x_i)$ and $\mathbb{P}(\mathbf{X} > \mathbf{x}) \leq \prod_{i=1}^d \mathbb{P}(X_i > x_i)$.

Moreover, we say that a distribution or a copula is PA, PRD, WPA, POD, NA, NRD, WNA, or NOD if the corresponding random vector is.

Note that the definition of PA does not require A and B to be disjoint, whereas the definition of NA requires this.

The relationship between the above notions is summarized below (see e.g., Chen, Embrechts, and Wang (2024a)):

 $PA \Longrightarrow WPA, PRD \Longrightarrow WPA, WPA \Longrightarrow POD,$

 $NA \Longrightarrow WNA$, $NRD \Longrightarrow WNA$, $WNA \Longrightarrow NOD$.

Within the class of multivariate normal distributions, the four concepts of positive dependence are equivalent, and each is equivalent to having nonnegative bivariate correlation coefficients; similarly, the four concepts of negative dependence are equivalent, and each is equivalent to having nonpositive bivariate correlation coefficients.

In the following, we use \mathfrak{D} to represent one of the following: PA, PRD, WPA, POD, NA, NRD, WNA, or NOD. Our question is whether these properties are properties purely based on copulas. It turns out that the checkerboard copula can help answer this question.

5.2. The checkerboard copula preserves dependence. We first present a self-consistency property of those negative dependence concepts in the spirit of Joag-Dev and Proschan (1983, Property P_6) for NA.

Lemma 2. If f_1, \ldots, f_d are increasing functions and \mathbf{X} satisfies \mathfrak{D} , then $(f_1(X_1), \ldots, f_d(X_d))$ also satisfies \mathfrak{D} .

Proof. We only show the result for the concepts of negative dependence, as the case of positive dependence is similar.

The self-consistency properties of NA and NOD are shown in Joag-Dev and Proschan (1983, Property P₆) and Lehmann (1966, Lemma 1), respectively. We will show the properties for NRD and WNA. Let $\mathbf{Y} = (f_1(X_1), \ldots, f_d(X_d))$.

1. Assume that **X** is NRD. Fix $i \in [d]$. Let g be a coordinatewise increasing function, and let $g' = g \circ (f_1, \ldots, f_{i-1}, f_{i+1}, \ldots, f_d)$. As a result, we have that g' is a coordinatewise increasing function and that $g(\mathbf{Y}_{-i}) = g'(\mathbf{X}_{-i})$. For any $y \in \mathbb{R}$, let $A_y = \{x : f_i(x) = y\}$. We have $\{Y_i = y\} = \{X_i \in A_y\}$. Therefore, $\mathbb{E}[g(\mathbf{Y}_{-i})|Y_i = y] = \mathbb{E}[g'(\mathbf{X}_{-i})|X_i \in A_y]$. Assume that $y_1 < y_2$. For any $x_1 \in A_{y_1}$ and $x_2 \in A_{y_2}$, we have $x_1 \leq x_2$; hence, $\mathbb{E}[g'(\mathbf{X}_{-i})|X_i = x_1] \ge \mathbb{E}[g'(\mathbf{X}_{-i})|X_i = x_2]$. Thus,

$$\mathbb{E}[g'(\mathbf{X}_{-i})|X_i \in A_{y_1}] = \mathbb{E}[\mathbb{E}[g'(\mathbf{X}_{-i})|X_i]|X_i \in A_{y_1}]$$

$$\geq \mathbb{E}[\mathbb{E}[g'(\mathbf{X}_{-i})|X_i]|X_i \in A_{y_2}] = \mathbb{E}[g'(\mathbf{X}_{-i})|X_i \in A_{y_2}],$$

which implies that $\mathbb{E}[g(\mathbf{Y}_{-i})|Y_i = y_1] \ge \mathbb{E}[g(\mathbf{Y}_{-i})|Y_i = y_2]$; hence, **Y** is NRD. 2. Assume that **X** is WNA. For $i \in [d]$, let $S \subseteq \mathbb{R}^{d-1}$ be a decreasing set, and let

$$S_i^f = \{(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_d) : (f_1(x_1), \dots, f_{i-1}(x_{i-1}), f_{i+1}(x_{i+1}), \dots, f_d(x_d)) \in S\}.$$

It is clear that $\{\mathbf{Y}_{-i} \in S\} = \{\mathbf{X}_{-i} \in S_i^f\}$. For any $\mathbf{x}_1 \leq \mathbf{x}_2$ and $\mathbf{x}_2 \in S_i^f$, we have $f_k(x_{1,k}) \leq f_k(x_{2,k})$ for all $k \in [d] \setminus \{i\}$. Furthermore, because S is decreasing, we have $\mathbf{x}_1 \in S_i^f$, which implies that S_i^f is a decreasing set. For any $y \in \mathbb{R}$ with $\mathbb{P}(Y_i \leq y) > 0$, let $x = \sup\{t \in \mathbb{R} : f_i(t) \leq y\}$. If $f_i(x) \leq y$, then we have $\{Y_i \leq y\} = \{X_i \leq x\}$ and $\mathbb{P}(X_i \leq x) > 0$. Therefore,

$$\mathbb{P}(\mathbf{Y}_{-i} \in S | Y_i \leqslant y) = \mathbb{P}\left(\mathbf{X}_{-i} \in S_i^f | X_i \leqslant x\right) \leqslant \mathbb{P}\left(\mathbf{X}_{-i} \in S_i^f\right) = \mathbb{P}(\mathbf{Y}_{-i} \in S),$$

which implies that **Y** is WNA. If $f_i(x) > y$, then we have $\{Y_i \leq y\} = \{X_i < x\}$ and $\mathbb{P}(X_i < x) > 0$. Therefore,

Copyright © by SIAM. Unauthorized reproduction of this article is prohibited.

$$\mathbb{P}(\mathbf{Y}_{-i} \in S, Y_i \leqslant y) = \mathbb{P}\left(\mathbf{X}_{-i} \in S_i^f, X_i < x\right)$$
$$= \lim_{t \uparrow x} \mathbb{P}\left(\mathbf{X}_{-i} \in S_i^f, X_i \leqslant t\right)$$
$$\leqslant \lim_{t \uparrow x} \mathbb{P}\left(\mathbf{X}_{-i} \in S_i^f\right) \mathbb{P}(X_i \leqslant t)$$
$$= \mathbb{P}\left(\mathbf{X}_{-i} \in S_i^f\right) \lim_{t \uparrow x} \mathbb{P}(X_i \leqslant t)$$
$$= \mathbb{P}\left(\mathbf{X}_{-i} \in S_i^f\right) \mathbb{P}(X_i < x) = \mathbb{P}(\mathbf{Y}_{-i} \in S)\mathbb{P}(Y_i \leqslant y),$$

which implies that $\mathbb{P}(\mathbf{Y}_{-i} \in S | Y_i \leq y) \leq \mathbb{P}(\mathbf{Y}_{-i} \in S)$ and that **Y** is WNA.

The following theorem demonstrates that the checkerboard copula of \mathbf{X} preserves the dependence information of \mathbf{X} .

Theorem 3. A random vector \mathbf{X} satisfies \mathfrak{D} if and only if it has a copula that satisfies \mathfrak{D} . Moreover, the copula can be chosen as the checkerboard copula $C_{\mathbf{X}}^{\perp}$.

Proof. The "if" part follows from Lemma 2 because, for $\mathbf{U} = (U_1, \ldots, U_d)$ following the copula of \mathbf{X} that satisfies \mathfrak{D} , we have that $(X_1, \ldots, X_d) = (F_1^{-1}(U_1), \ldots, F_d^{-1}(U_d))$ and that F_i^{-1} is increasing for all $i \in [d]$.

Now we show the "only if" part. Let $\mathbf{U} = (U_1, \ldots, U_d)$ be the random vector given by (4) with $\mathbf{V} = (V_1, \ldots, V_d) \sim \mathbf{U}([0, 1]^d)$ independent of \mathbf{X} . Hence, we have that $\mathbf{U} \sim C_{\mathbf{X}}^{\perp}$ and that $C_{\mathbf{X}}^{\perp}$ is a copula of \mathbf{X} . Note that for any $i \in [d]$, given V_i , we have that U_i is an increasing function of X_i . Hence, by Lemma 2, \mathbf{X} satisfies \mathfrak{D} implies that $\mathbf{U}|\mathbf{V}$ also satisfies \mathfrak{D} .

Assume that **X** is NA. For any given pair of disjoint subsets A, B of [d] and any given functions f and g both increasing or decreasing coordinatewise, we have

$$\operatorname{Cov}(f(\mathbf{U}_A), g(\mathbf{U}_B)) = \mathbb{E}[\operatorname{Cov}(f(\mathbf{U}_A), g(\mathbf{U}_B) | \mathbf{V})] + \operatorname{Cov}(\mathbb{E}[f(\mathbf{U}_A) | \mathbf{V}], \mathbb{E}[g(\mathbf{U}_B) | \mathbf{V}])$$

$$\leq 0 + \operatorname{Cov}(\mathbb{E}[f(\mathbf{U}_A) | \mathbf{V}_A], \mathbb{E}[g(\mathbf{U}_B) | \mathbf{V}_B]) = 0,$$

where the inequality follows from the fact that $\mathbf{U}|\mathbf{V}$ is NA and the last equality follows from the independence between \mathbf{V}_A and \mathbf{V}_B . Hence, \mathbf{U} is NA.

Assume that **X** is NRD. For any fixed *i* and *k*, by (4), there exist *x* and *v* such that $\{U_i = k\} = \{X_i = x, V_i = v\}$. Then for any coordinatewise increasing function *g*, by the independence between V_i and (X_i, \mathbf{U}_{-i}) , we have

$$\mathbb{E}[g(\mathbf{U}_{-i})|U_i = k] = \mathbb{E}[g(\mathbf{U}_{-i})|X_i = x, V_i = v] = \mathbb{E}[g(\mathbf{U}_{-i})|X_i = x].$$

Because \mathbf{U}_{-i} is a function of \mathbf{X}_{-i} and \mathbf{V}_{-i} , we can let *h* be the function such that $g(\mathbf{U}_{-i}) = h(\mathbf{X}_{-i}, \mathbf{V}_{-i})$. Then due to the independence between \mathbf{V}_{-i} and \mathbf{X} ,

$$\mathbb{E}[g(\mathbf{U}_{-i})|X_i = x] = \mathbb{E}[h(\mathbf{X}_{-i}, \mathbf{V}_{-i})|X_i = x] = \int_{[0,1]^{d-1}} \mathbb{E}[h(\mathbf{X}_{-i}, \mathbf{v}_{-i})|X_i = x] d\mathbf{v}_{-i},$$

where $\mathbf{v}_{-i} = (v_1, \ldots, v_{i-1}, v_{i+1}, \ldots, v_d)$. Therefore, for any $k_1 \leq k_2$, there exist x_1 and x_2 such that

Copyright (C) by SIAM. Unauthorized reproduction of this article is prohibited.

$$\mathbb{E}[g(\mathbf{U}_{-i})|U_{i} = k_{1}] = \int_{[0,1]^{d-1}} \mathbb{E}[h(\mathbf{X}_{-i}, \mathbf{v}_{-i})|X_{i} = x_{1}] d\mathbf{v}_{-i},$$
$$\mathbb{E}[g(\mathbf{U}_{-i})|U_{i} = k_{2}] = \int_{[0,1]^{d-1}} \mathbb{E}[h(\mathbf{X}_{-i}, \mathbf{v}_{-i})|X_{i} = x_{2}] d\mathbf{v}_{-i}.$$

In addition, by (4), we must have $x_1 \leq x_2$. Note that given \mathbf{v}_{-i} , $h(\mathbf{X}_{-i}, \mathbf{v}_{-i})$ is a coordinatewise increasing function of \mathbf{X}_{-i} . Hence, we have $\mathbb{E}[h(\mathbf{X}_{-i}, \mathbf{v}_{-i})|X_i = x_1] \ge \mathbb{E}[h(\mathbf{X}_{-i}, \mathbf{v}_{-i})|X_i = x_2]$ for any \mathbf{v}_{-i} . Therefore, $\mathbb{E}[g(\mathbf{U}_{-i})|U_i = k_1] \ge \mathbb{E}[g(\mathbf{U}_{-i})|U_i = k_2]$, and \mathbf{U} is NRD.

Assume that **X** is WNA. For any $i \in [d]$, decreasing set $S \subseteq \mathbb{R}^{d-1}$, and $x \in \mathbb{R}$ with $\mathbb{P}(U_i \leq x) > 0$,

$$\mathbb{P}(\mathbf{U}_{-i} \in S, U_i \leq x) = \mathbb{E}[\mathbb{P}(\mathbf{U}_{-i} \in S, U_i \leq x \mid \mathbf{V})]$$

$$\leq \mathbb{E}[\mathbb{P}(\mathbf{U}_{-i} \in S \mid \mathbf{V}_{-i})\mathbb{P}(U_i \leq x \mid V_i)]$$

$$= \mathbb{E}[\mathbb{P}(\mathbf{U}_{-i} \in S \mid \mathbf{V}_{-i})]\mathbb{E}[\mathbb{P}(U_i \leq x \mid V_i)]$$

$$= \mathbb{P}(\mathbf{U}_{-i} \in S)\mathbb{P}(U_i \leq x).$$

Hence, \mathbf{U} is WNA.

Assume that **X** is NOD. For any $t_1, \ldots, t_d \in \mathbb{R}$, we have

$$\mathbb{P}(U_1 \leqslant t_1, \dots, U_d \leqslant t_d) = \mathbb{E}[\mathbb{P}(U_1 \leqslant t_1, \dots, U_d \leqslant t_d | V_1, \dots, V_d)]$$

$$\leqslant \mathbb{E}[\mathbb{P}(U_1 \leqslant t_1 | V_1) \cdots \mathbb{P}(U_d \leqslant t_d | V_d)]$$

$$= \mathbb{E}[\mathbb{P}(U_1 \leqslant t_1 | V_1)] \cdots \mathbb{E}[\mathbb{P}(U_d \leqslant t_d | V_d)]$$

$$= \mathbb{P}(U_1 \leqslant t_1) \cdots \mathbb{P}(U_d \leqslant t_d).$$

Similarly, we can show that

$$\mathbb{P}(U_1 > t_1, \dots, U_d > t_d) \leqslant \mathbb{P}(U_1 > t_1) \cdots \mathbb{P}(U_d > t_d).$$

Hence, \mathbf{U} is NOD.

In conclusion, if **X** satisfies \mathfrak{D} , then **U** satisfies \mathfrak{D} , where \mathfrak{D} is one of the four concepts of negative dependence.

To show the case of positive dependence, we follow a similar route. We take the same **U** as above. Assume that **X** is PA. Because $U_i|V_i$ is an increasing function of X_i , by Lemma 2, **U**|**V** is also PA. Thus, for any given pair of subsets A, B of [d] and any given functions f and g both coordinatewise increasing or decreasing, we have

$$\operatorname{Cov}(f(\mathbf{U}_A), g(\mathbf{U}_B)) = \mathbb{E}[\operatorname{Cov}(f(\mathbf{U}_A), g(\mathbf{U}_B) | \mathbf{V})] + \operatorname{Cov}(\mathbb{E}[f(\mathbf{U}_A) | \mathbf{V}], \mathbb{E}[g(\mathbf{U}_B) | \mathbf{V}])$$

$$\geq \operatorname{Cov}(\mathbb{E}[f(\mathbf{U}_A) | \mathbf{V}_A], \mathbb{E}[g(\mathbf{U}_B) | \mathbf{V}_B]).$$

Moreover, given \mathbf{X} , U_i is an increasing function of V_i . Hence, $\mathbb{E}[f(\mathbf{U}_A)|\mathbf{V}_A]$ and $\mathbb{E}[f(\mathbf{U}_B)|\mathbf{V}_B]$ are coordinatewise increasing (or decreasing) with respect to \mathbf{V}_A and \mathbf{V}_B , respectively, if f and g are both coordinatewise increasing (or decreasing). Because \mathbf{V} is PA, we have

$$\operatorname{Cov}\left(\mathbb{E}[f(\mathbf{U}_A)|\mathbf{V}_A], \mathbb{E}[g(\mathbf{U}_B)|\mathbf{V}_B]\right) \ge 0,$$

Copyright (c) by SIAM. Unauthorized reproduction of this article is prohibited.

implying that **U** is PA. The proofs for other positive dependence concepts are similar. A partial proof for POD can also be found in Durante et al. (2015, Proposition 2.3).

Genest and Nešlehová (2007) also show the dependence preservation results for positive orthant dependence, positive likelihood ratio dependence, and tail dependence in the bivariate case.

Remark 2. We can clearly see from Theorem 3 that $C_{\mathbf{X}}^{\perp} = \Pi$ (the independence copula) is the only independent copula in $\mathcal{C}_{\mathbf{X}}$ when \mathbf{X} is independent. This fact is used as the basis of the independence test in Genest et al. (2019).

6. Two consequences of Theorem **3**. We provide two applications in this section to highlight the usefulness of Theorem 3.

6.1. Diversification penalty. For random variables X and Y, let $X \ge_{\text{st}} Y$ represent $\mathbb{P}(X > x) \ge \mathbb{P}(Y > x)$ for all $x \in \mathbb{R}$; this is called the stochastic order. Chen, Embrechts, and Wang (2024a, 2024b) studied the problem of diversification penalty, that is, whether

(5)
$$X \leq_{\text{st}} \sum_{i=1}^{d} \theta_i X_i$$
 for all $(\theta_1, \dots, \theta_d) \in \Delta_d$, where X, X_1, \dots, X_d are identically distributed,

holds under certain marginal distributions and dependence structures. Here Δ_d is the standard simplex defined by $\Delta_d = \{(\theta_1, \dots, \theta_d) \in [0, 1]^d : \theta_1 + \dots + \theta_d = 1\}$. When X is interpreted as a loss, (5) intuitively means that the nondiversified portfolio X is less dangerous than the diversified portfolio $\sum_{i=1}^{d} \theta_i X_i$. This seems counterintuitive at first glance, but it indeed happens in the model of Chen, Embrechts, and Wang (2024a), where X has infinite mean.

Define the set, for some dependence concept \mathfrak{D} in section 5.1,

$$\mathcal{F}_{\mathfrak{D}} = \{ \text{distribution of } X : (5) \text{ holds for all } (X_1, \dots, X_d) \text{ that satisfy } \mathfrak{D} \}.$$

Chen, Embrechts, and Wang (2024a) showed that the Pareto(1) distribution belongs to \mathcal{F}_{WNA} and hence also to \mathcal{F}_{NA} , \mathcal{F}_{NRD} , and \mathcal{F}_{IN} , where IN stands for independence. Moreover, Chen, Embrechts, and Wang (2024b, Proposition 1) showed that $\mathcal{F}_{\mathfrak{D}}$ for \mathfrak{D} being WNA, NA, or IN is closed under strictly increasing convex transforms on the random variables. Our next result, which relies on our Theorem 3, addresses non–strictly increasing f and other notions of dependence, thus generalizing the above result.

Proposition 2. Each of $\mathcal{F}_{\mathfrak{D}}$ is closed under increasing convex transforms on the random variable.

Proof. Below we first show that each of $\mathcal{F}_{\mathfrak{D}}$ is closed under strictly increasing convex transforms on the random variable; that is, if the distribution of X is in $\mathcal{F}_{\mathfrak{D}}$, then so is the distribution of f(X) for a strictly increasing convex f. Assume that $F \in \mathcal{F}_{\mathfrak{D}}$, X follows F and that Y = f(X), where f is strictly increasing and convex. Because f is strictly increasing, if (Y_1, \ldots, Y_d) satisfies \mathfrak{D} , then so does (X_1, \ldots, X_d) , where $X_i = f^{-1}(Y_i)$ for $i \in [d]$, by Lemma 2. Because each of X, X_1, \ldots, X_d has a distribution $F \in \mathcal{F}_{\mathfrak{D}}$, we have $X \leq_{\mathrm{st}} \sum_{i=1}^d \theta_i X_i$, and this gives, using the convexity of f,

(6)
$$Y = f(X) \leqslant_{\text{st}} f\left(\sum_{i=1}^{d} \theta_i X_i\right) \leqslant \sum_{i=1}^{d} \theta_i f(X_i) = \sum_{i=1}^{d} \theta_i Y_i.$$

To address the case that f is not strictly increasing, Theorem 3 allows us to find the above (X_1, \ldots, X_d) that satisfies \mathfrak{D} and such that $Y_i = f(X_i)$ for $i \in [d]$. In particular, using Theorem 3, we can construct (U_1, \ldots, U_d) , which follows the checkerboard copula of (Y_1, \ldots, Y_d) and satisfies \mathfrak{D} such that

$$(Y_1,\ldots,Y_d)=(f\circ g(U_1),\ldots,f\circ g(U_d)),$$

where g is the quantile function of X and $f \circ g$ is the quantile function of Y. Setting $(X_1, \ldots, X_d) = (g(U_1), \ldots, g(U_d))$, we get that (X_1, \ldots, X_d) satisfies \mathfrak{D} , and this leads to (6).

6.2. Induced order statistics. Here we demonstrate another application of Theorem 3 in characterizing the distribution of induced order statistics. Consider N independent and identically distributed bivariate random vectors

$$\begin{pmatrix} \xi_1\\ \eta_1 \end{pmatrix}, \begin{pmatrix} \xi_2\\ \eta_2 \end{pmatrix}, \dots, \begin{pmatrix} \xi_N\\ \eta_N \end{pmatrix}.$$

Note that for $i \neq j$, (ξ_i, η_i) and (ξ_j, η_j) are independent and identically distributed but that ξ_i and η_i may be correlated and have different marginal distributions. We rank these bivariate vectors according to their first components, ξ_i ,

(7)
$$\begin{pmatrix} \xi_{1:N} \\ \eta_{[1:N]} \end{pmatrix}, \begin{pmatrix} \xi_{2:N} \\ \eta_{[2:N]} \end{pmatrix}, \dots, \begin{pmatrix} \xi_{N:N} \\ \eta_{[N:N]} \end{pmatrix},$$

where $\xi_{1:N} \leq \xi_{2:N} \leq \cdots \leq \xi_{N:N}$ are the order statistics of $\xi_1, \xi_2, \ldots, \xi_N$. The notation $\eta_{[i:N]}$ represents the *i*th *induced order statistic* (Bhattacharya (1974)), where the order is induced by another variable ξ_i . The induced order statistics $\eta_{[1:N]}, \ldots, \eta_{[N:N]}$ are also referred to as *concomitants* of the order statistics $\xi_{1:N}, \ldots, \xi_{N:N}$ (David (1973)).

In the context of constructing impact portfolios, Lo et al. (2024) investigated the joint distribution of $(\eta_{[1:N]}, \ldots, \eta_{[N:N]})$. In particular, they proved a representation theorem for the joint distribution of $(\eta_{[1:N]}, \ldots, \eta_{[N:N]})$ using the copula of (ξ_i, η_i) . Furthermore, they demonstrated that if ξ_i is not continuously distributed, then the representation theorem holds if and only if the copula of (ξ_i, η_i) is chosen as the (bivariate) checkerboard copula in this paper. This reveals a potential application of the checkerboard copula in portfolio construction.

Lo et al. (2024) also showed that the rank of the odd-order moments of induced order statistics relies on the copula of (ξ_i, η_i) . Assume that C is a copula of (ξ_i, η_i) . Lo et al. (2024, Theorem EC.5) proved that for any k = 0, 1, ..., if C is PRD, then we have

(8)
$$\mathbb{E}\left(\eta_{[1:N]}^{2k+1}\right) \leqslant \mathbb{E}\left(\eta_{[2:N]}^{2k+1}\right) \leqslant \dots \leqslant \mathbb{E}\left(\eta_{[N:N]}^{2k+1}\right),$$

and if C is NRD, then we have

(9)
$$\mathbb{E}\left(\eta_{[1:N]}^{2k+1}\right) \ge \mathbb{E}\left(\eta_{[2:N]}^{2k+1}\right) \ge \dots \ge \mathbb{E}\left(\eta_{[N:N]}^{2k+1}\right).$$

In particular, the copula C can be chosen as the checkerboard copula. Therefore, using Theorem 3, we directly obtain the following result.

Proposition 3. For any k = 0, 1, ..., (8) holds if (ξ_i, η_i) is PRD, and (9) holds if (ξ_i, η_i) is NRD.

The difference between Proposition 3 and Lo et al. (2024, Theorem EC.5) is that the latter imposes the dependence assumption (PRD or NRD) on the copula of (ξ_i, η_i) , while the former imposes a more natural assumption on the random vector (ξ_i, η_i) directly, which is only possible due to our Theorem 3.

7. Applications in corisk measures and portfolio selection. In this section, we use both numerical and empirical experiments to show that the choice of copula impacts the calculation of corisk measures when the marginal distributions are not continuous. In particular, we consider marginal ES as defined in (1), which is discussed in our third motivating question in section 1. Section 7.1 presents a numerical experiment to show that different copula choices can lead to varying marginal ES results. Section 7.2 uses real data from the U.S. stock market to illustrate how the choice of copula affects the performance of the portfolio with minimum marginal ES. Our results not only highlight the importance of copula selection in the computation of corisk measures in financial practice but also show that the checkerboard copula is often the most convenient and natural choice that can produce reliable results.

7.1. Numerical experiment. Consider a bivariate normal distribution with marginals $N(0, \sigma^2)$ and correlation r. Denote this bivariate distribution by F_r . We choose this distribution because it is well known that for $(X_1, X_2) \sim F_r$, the marginal ES of X_2 given X_1 at level $p \in (0, 1)$ can be explicitly computed as

(10)
$$\rho(X_2|X_1) = \mathbb{E}[X_2|X_1 > \Phi^{-1}(p)] = \frac{r\sigma}{1-p}\varphi\left(\Phi^{-1}(p)\right),$$

where φ and Φ are the density and distribution functions of N(0,1), respectively.

Now we conduct a numerical experiment based on this bivariate normal distribution. We draw 1,000 bivariate random vectors, $(X_1^{(1)}, X_2^{(1)}), \ldots, (X_1^{(1,000)}, X_2^{(1,000)})$, from F_r . These random values are then rounded to one decimal place to estimate an empirical bivariate distribution, denoted by \hat{F} . Thus, \hat{F} is a discrete bivariate distribution on the discrete grid caused by rounding. Next, assuming that $(\hat{X}_1, \hat{X}_2) \sim \hat{F}$, we compute the marginal ES of \hat{X}_2 given \hat{X}_1 at level $p \in (0, 1)$, which is $\rho(\hat{X}_2|\hat{X}_1)$ defined as (1). When calculating the marginal ES, the following two copulas of (\hat{X}_1, \hat{X}_2) given by (4) are considered:

- (i) Let $(V_1, V_2) \sim U([0, 1]^2)$ be independent of (X_1, X_2) ; that is, use the checkerboard copula C^{\perp} .
- (ii) Let $V_2 \sim U[0,1]$ be independent of \hat{X}_2 . In addition, given \hat{X}_2 , let V_2 and \hat{X}_1 be comonotonic. Let $V_1 \sim U[0,1]$ be independent of \hat{X}_1 , \hat{X}_2 , and V_2 . We denote this copula by C^+ .

Therefore, we obtain different values of $\rho(\hat{X}_2|\hat{X}_1)$ using the two different copulas.

Given (r, p), we run the simulation procedure described above 10,000 times and calculate the marginal ES under both copulas for each run. We choose $\sigma = 10$. Table 1 shows the marginal ES given by (10) (normal formula), the average marginal ES value, and the mean squared error (MSE)—the mean squared difference between $\rho(\hat{X}_2|\hat{X}_1)$ and the value in (10) across the 10,000 runs for each of the two copulas.

	p = 0.9				p = 0.95		p = 0.975			
ρ	0.2	0.3	0.4	0.2	0.3	0.4	0.2	0.3	0.4	
Normal formula	3.510	5.265	7.020	4.125	6.188	8.251	4.676	7.013	9.351	
Average with C^{\perp}	3.502	5.257	7.001	4.116	6.171	8.234	4.688	6.992	9.330	
Average with C^+	3.985	5.730	7.455	4.687	6.726	8.767	5.338	7.615	9.934	
MSE with C^{\perp}	0.940	0.920	0.891	1.914	1.796	1.673	3.725	3.636	3.329	
MSE with C^+	1.225	1.192	1.131	2.348	2.173	2.020	4.357	4.187	3.796	

 Table 1

 Simulation results for computing marginal ES under two copulas.

Table 2																
Marginal	ES	of	the	loss	of	five	individual	stocks	given	the	loss	of	the	$S \mathscr{C} P$	500	Index.

	Microsoft	Apple	Google	Nvidia	Amazon
C^{\perp} C^+	3.518% 4.404%	3.545% 4.752%	3.455% 4.523%	5.043% 6.686%	3.712% 4.998%
Empirical distribution	3.845%	3.905%	3.735%	5.456%	4.054%

Table 1 illustrates that the choice of copula affects the calculation of the marginal ES. Under our setup, the marginal ES computed using the checkerboard copula is, on average, closer to the result obtained using the normal formula (10). This demonstrates that the checkerboard copula is a good candidate for computing corisk measures for noncontinuous random variables.

7.2. Empirical study. To further demonstrate the importance of copula selection when computing corisk measures in financial practice, we use real stock data to calculate the marginal ES. We obtain daily returns of the S&P 500 Index and five widely traded U.S. stocks—Microsoft, Apple, Google, Nvidia, and Amazon—from 2005 to 2023.² To make the distribution discrete, we classify market conditions into five groups based on the daily returns of the S&P 500 Index: $(-\infty, -3\%]$, (-3%, -1%], (-1%, +1%], (+1%, +3%], and $(+3\%, +\infty)$. These five conditions represent very bad, bad, fair, good, and very good, with corresponding values of -2, -1, 0, +1, and +2, respectively.

Next, for each individual stock, we use the negative values of its daily returns along with these market condition values over the entire period to estimate an empirical bivariate distribution. Then based on the empirical bivariate distribution, the two copulas described in section 7.1 are applied to calculate the marginal ES of the stock's loss given the market condition. We also present results computed using the empirical bivariate distributions of daily stock returns and S&P 500 Index returns directly, without classifying the market conditions.

Table 2 shows the marginal ES of the five stocks under the two copulas, along with the result computed directly from the empirical distributions. The marginal ES values differ across the three methods, and the results from the checkerboard copula, C^{\perp} , are lower than those from the alternative copula, C^{+} .

²Our data come from the Center for Research in Security Prices (CRSP). We use data starting in 2005 because Google became publicly traded in August 2004.

THE CHECKERBOARD COPULA AND DEPENDENCE CONCEPTS



Figure 1. Cumulative values of the minMES portfolios.

Table 3

Performance metrics (average annual returns, standard deviation of annual returns, and Sharpe ratio) for the minMES portfolios. The Sharpe ratio is calculated assuming a risk-free rate of 3%.

Metrics	Average return	Standard deviation	Sharpe ratio
C^{\perp}	22.34%	29.89%	0.6470
C^+	19.73%	27.65%	0.6050
Empirical distribution	20.11%	29.92%	0.5721

Different choices of copulas can also result in varying financial performance for the portfolio with minimum marginal ES (minMES). To demonstrate this, we construct minMES portfolios for the five stocks as follows. At the beginning of year t, we determine the weights of the five stocks that minimize the marginal ES of the portfolio's loss given the market condition, using data from year t - 1. The optimization is subject to the constraints that all weights must be nonnegative and sum to 1. These optimal weights are then held throughout year t.

Figure 1 and Table 3 show the cumulative portfolio values and performance metrics of this minMES strategy for p = 0.975 under the three methods over the entire sample period, respectively. We find that the choice of copula significantly impacts the financial performance of the minMES portfolio. Furthermore, in our empirical study, the checkerboard copula generally achieves better performance, demonstrating that it is a convenient and effective choice for producing reliable results for the considered dataset. Certainly, we do not claim that this advantage is profitable in general, which requires comprehensive empirical analysis.

8. Conclusion. We discussed the choice of copula when the marginal distributions are not necessarily continuous. Among all the choices of copulas for a given random vector, the checkerboard copula is the most convenient and natural selection in applications such as simulating from the copula, stressing the distribution, and computing a corisk measure. It is shown that the checkerboard copula is the most unbiased choice in the sense that it has the largest Shannon entropy among all possible copulas for a given random vector. Moreover, the checkerboard copula can preserve the dependence information of the underlying random vector. This preservation property is applied to identify suitable distributions in the context of diversification penalty studied by Chen, Embrechts, and Wang (2024a, 2024b) and to determine the ranks of the moments of induced order statistics in the context of impact portfolios studied by Lo et al. (2024). Finally, our results indicate that the choice of copula significantly affects the calculation of corisk measures when the marginal distributions are not continuous. Through numerical experiments and empirical studies, we find that the checkerboard copula can produce reliable results when computing marginal ES and demonstrate strong performance when constructing minMES portfolios.

REFERENCES

- T. ADRIAN AND M. K. BRUNNERMEIER (2016), CoVaR, Amer. Econ. Rev., 106, pp. 1705–1741, https://doi.org/ 10.1257/aer.20120555.
- K. ALAM AND K. M. L. SAXENA (1981), Positive dependence in multivariate distributions, Comm. Statist. Theory Methods, 10, pp. 1183–1196, https://doi.org/10.1080/03610928108828102.
- Y. BENJAMINI AND D. YEKUTIELI (2001), The control of the false discovery rate in multiple testing under dependency, Ann. Stat., 29, pp. 1165–1188, https://doi.org/10.1214/aos/1013699998.
- P. K. BHATTACHARYA (1974), Convergence of sample paths of normalized sums of induced order statistics, Ann. Stat., 2, pp. 1034–1039, https://doi.org/10.1214/aos/1176342823.
- H. W. BLOCK, T. H. SAVITS, AND M. SHAKED (1982), Some concepts of negative dependence, Ann. Probab., 10, pp. 765–772, https://doi.org/10.1214/aop/1176993784.
- H. W. BLOCK, T. H. SAVITS, AND M. SHAKED (1985), A concept of negative dependence using stochastic ordering, Statist. Probab. Lett., 3, pp. 81–86, https://doi.org/10.1016/0167-7152(85)90029-X.
- H. CARLEY (2002), Maximum and minimum extensions of finite subcopulas, Comm. Statist. Theory Methods, 31, pp. 2151–2166, https://doi.org/10.1081/STA-120017218.
- Y. CHEN, P. EMBRECHTS, AND R. WANG (2024a), Technical note—An unexpected stochastic dominance: Pareto distributions, dependence, and diversification, Oper. Res., https://doi.org/10.1287/opre.2022.0505.
- Y. CHEN, P. EMBRECHTS, AND R. WANG (2024b), Risk Exchange under Infinite-Mean Pareto Models, preprint, arXiv:2403.20171.
- T. M. COVER AND J. A. THOMAS (1991), Elements of Information Theory, John Wiley, New York.
- I. CSISZÁR (1975), I-divergence geometry of probability distributions and minimization problems, Ann. Probab., 3, pp. 146–158, https://www.jstor.org/stable/2959270.
- H. A. DAVID AND H. N. NAGARAJA (1998), Concomitants of order statistics, Handbook of Statist., 16, pp. 487–513, https://doi.org/10.1016/S0169-7161(98)16020-0.
- E. DE AMO, M. D. CARRILLO, F. DURANTE, AND J. F. SÁNCHEZ (2017), Extensions of subcopulas, J. Math. Anal. Appl., 452, pp. 1–15, https://doi.org/10.1016/j.jmaa.2017.02.061.
- M. DENUIT AND P. LAMBERT (2005), Constraints on concordance measures in bivariate discrete data, J. Multivariate Anal., 93, pp. 40–57, https://doi.org/10.1016/j.jmva.2004.01.004.
- F. DURANTE, J. FERNÁNDEZ-SÁNCHEZ, J. J. QUESADA-MOLINA, AND M. UBEDA-FLORES (2015), Convergence results for patchwork copulas, Eur. J. Oper. Res., 247, pp. 525–531, https://doi.org/10.1016/j.ejor. 2015.06.028.
- J. D. ESARY, F. PROSCHAN, AND D. W. WALKUP (1967), Association of random variables, with applications, Ann. Math. Stat., 38, pp. 1466–1474, https://doi.org/10.1214/aoms/1177698701.
- O. P. FAUGERAS (2017), Inference for copula modeling of discrete data: A cautionary tale and some facts, Depend. Model., 5, pp. 121–132, https://doi.org/10.1515/demo-2017-0008.
- T. S. FERGUSON (1967), Mathematical Statistics: A Decision Theoretic Approach, Academic Press, New York.
- G. GEENENS (2020), Copula modeling for discrete random vectors, Depend. Model., 8, pp. 417–440, https:// doi.org/10.1515/demo-2020-0022.
- C. GENEST AND J. NEŠLEHOVÁ (2007), A primer on copulas for count data, Astin Bull., 37, pp. 475–515, https://doi.org/10.2143/AST.37.2.2024077.
- C. GENEST, J. G. NEŠLEHOVÁ, AND B. RÉMILLARD (2013), On the estimation of Spearman's rho and related tests of independence for possibly discontinuous multivariate data, J. Multivariate Anal., 117, pp. 214–228, https://doi.org/10.1016/j.jmva.2013.02.007.

THE CHECKERBOARD COPULA AND DEPENDENCE CONCEPTS

- C. GENEST, J. G. NEŠLEHOVÁ, AND B. RÉMILLARD (2017), Asymptotic behavior of the empirical multilinear copula process under broad conditions, J. Multivariate Anal., 159, pp. 82–110, https://doi.org/10.1016/ j.jmva.2017.04.002.
- C. GENEST, J. G. NEŠLEHOVÁ, B. RÉMILLARD, AND O. A. MURPHY (2019), Testing for independence in arbitrary distributions, Biometrika, 106, pp. 47–68, https://doi.org/10.1093/biomet/asy059.
- E. T. JAYNES (1957), Information theory and statistical mechanics, Phys. Rev., 106, pp. 620–628, https:// doi.org/10.1103/PhysRev.106.620.
- K. JOAG-DEV AND F. PROSCHAN (1983), Negative association of random variables with applications, Ann. Stat., 11, pp. 286–295, https://doi.org/10.1214/aos/1176346079.
- H. JOE (2015), Dependence Modeling with Copulas, Chapman and Hall, London.
- G. KOLIANDER, G. PICHLER, E. RIEGLER, AND F. HLAWATSCH (2016), Entropy and source coding for integerdimensional singular random variables, IEEE Trans. Inform. Theory, 62, pp. 6124–6154, https://doi.org/ 10.1109/TIT.2016.2604248.
- V. KUZMENKO, R. SALAM, AND S. URYASEV (2020), Checkerboard copula defined by sums of random variables, Depend. Model., 8, pp. 70–92, https://doi.org/10.1515/demo-2020-0004.
- E. L. LEHMANN (1966), Some concepts of dependence, Ann. Math. Stat., 37, pp. 1137–1153, https://doi.org/ 10.1214/aoms/1177699260.
- D. X. LI (2000), On default correlation: A copula function approach, J. Fixed Income, 9, pp. 43–54, https:// doi.org/10.3905/jfi.2000.319253.
- A. W. LO, L. WU, R. ZHANG, AND C. ZHAO (2024), Optimal impact portfolios with general dependence and marginals, Oper. Res., 72, pp. 1775–1789, https://doi.org/10.1287/opre.2023.0400.
- A. W. MARSHALL (1996), Copulas, marginals, and joint distributions, in Distributions with Fixed Marginals and Related Topics, L, Rüschendorf, B. Schweizer, and M. D. Taylor, eds., Institute of Mathematical Statistics Lecture Notes-Monograph Series 28, Institute of Mathematical Statistics, Hayward, CA, pp. 213–222.
- A. J. MCNEIL, R. FREY, AND P. EMBRECHTS (2015), Quantitative Risk Management: Concepts, Techniques and Tools, rev. ed., Princeton University Press, Princeton, NJ.
- M. MESFIOUI AND J. F. QUESSY (2010), Concordance measures for multivariate non-continuous random vectors, J. Multivariate Anal., 101, pp. 2398–2410, https://doi.org/10.1016/j.jmva.2010.06.011.
- P. MILLOSSOVICH, A. TSANAKAS, AND R. WANG (2024), A theory of multivariate stress testing, Eur. J. Oper. Res., 318, pp. 851–866, https://doi.org/10.1016/j.ejor.2024.06.002.
- D. S. MOORE AND M. C. SPRUILL (1975), Unified large-sample theory of general chi-squared statistics for tests of fit, Ann. Stat., 3, pp. 599–616, https://doi.org/10.1214/aos/1176343125.
- R. NELSEN (2006), An Introduction to Copulas, 2nd ed., Springer-Verlag, Berlin.
- J. NEŠLEHOVÁ (2004), Dependence of Non-Continuous Random Variables, Doctoral dissertation, Universität Oldenburg, Oldenburg, Germany.
- J. NEŠLEHOVÁ (2007), On rank correlation measures for non-continuous random variables, J. Multivariate Anal., 98, pp. 544–567, https://doi.org/10.1016/j.jmva.2005.11.007.
- E. PERRONE, L. SOLUS, AND C. UHLER (2019), Geometry of discrete copulas, J. Multivariate Anal., 172, pp. 162–179, https://doi.org/10.1016/j.jmva.2019.01.014.
- J. PIANTADOSI, P. HOWLETT, AND J. BORWEIN (2012), Copulas with maximum entropy, Optim. Lett., 6, pp. 99–125, https://doi.org/10.1007/s11590-010-0254-2.
- L. RÜSCHENDORF (1981), Stochastically ordered distributions and monotonicity of the OC-function of sequential probability ratio tests, Statistics, 12, pp. 327–338, https://doi.org/10.1080/02331888108801593.
- L. RÜSCHENDORF (2009), On the distributional transform, Sklar's theorem, and the empirical copula process, J. Statist. Plann. Inference, 139, pp. 3921–3927, https://doi.org/10.1016/j.jspi.2009.05.030.
- L. RÜSCHENDORF (2013), Mathematical Risk Analysis: Dependence, Risk Bounds, Optimal Allocations and Portfolios, Springer-Verlag, Berlin.
- R. WANG AND R. ZITIKIS (2021), An axiomatic foundation for the expected shortfall, Manage. Sci., 67, pp. 1413–1429, https://doi.org/10.1287/mnsc.2020.3617.