

## 差分方法II, 作业3

交作业时间: 2023/04/28

1. For the scalar conservation law, show that LxF (Lax-Friedrichs), LLF (Local Lax-Friedrichs), and Engquist-Osher fluxes are all E-fluxes.
2. For a monotone flux we have that  $\mathcal{F}(\uparrow, \downarrow)$ . Show that a monotone flux is an E-flux.
3. In the MUSCL scheme, if the following conditions hold

$$\begin{aligned} \min(Q_{i-1}^n, Q_i^n) &\leq Q_{i-1/2}^{+,n+1/2} \leq \max(Q_{i-1}^n, Q_i^n), \\ \min(Q_i^n, Q_{i+1}^n) &\leq Q_{i+1/2}^{-,n+1/2} \leq \max(Q_i^n, Q_{i+1}^n), \end{aligned}$$

and the flux is monotone, then the MUSCL scheme is TVD-stable.

4. Consider the modified  $s$ -stage SSPRK for  $q_t = -\mathcal{L}(q)$ :

$$\begin{cases} Q^{(i)} = v_i Q^n + \sum_{j=1}^s [\alpha_{ij} Q^{(j)} + k \beta_{ij} \mathcal{L}(Q^{(j)})] & i = 1, \dots, s+1, \\ Q^{n+1} = Q^{(s+1)}. \end{cases}$$

Define  $\alpha, \beta \in \mathbb{R}^{(s+1) \times (s+1)}$  as

$$\alpha = \begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1s} & 0 \\ \alpha_{21} & \cdots & \alpha_{2s} & 0 \\ \vdots & & \vdots & \vdots \\ \alpha_{s+1,1} & \cdots & \alpha_{s+1,s} & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} \beta_{11} & \cdots & \beta_{1s} & 0 \\ \beta_{21} & \cdots & \beta_{2s} & 0 \\ \vdots & & \vdots & \vdots \\ \beta_{s+1,1} & \cdots & \beta_{s+1,s} & 0 \end{pmatrix}.$$

Show that the above SSPRK can be recast into the  $s$ -stage Runge-Kutta method:

$$\begin{cases} Q^{(i)} = Q^n + k \sum_{j=1}^s a_{ij} \mathcal{L}(Q^{(j)}) & i = 1, \dots, s, \\ Q^{n+1} = Q^n + k \sum_{i=1}^s b_i \mathcal{L}(Q^{(i)}), \end{cases}$$

under the following condition:

$$(\mathbf{I} - \alpha)^{-1} \beta = \begin{pmatrix} A & 0 \\ b^T & 0 \end{pmatrix}.$$

Here,  $A = (a_{ij}) \in \mathbb{R}^{s \times s}$ ,  $b = (b_1, \dots, b_s)^T \in \mathbb{R}^s$ .

5. Consider the WENO reconstruction

$$v_{i+1/2} = \sum_{r=0}^{m-1} \omega_r v_{i+1/2}^{(r)},$$

where the weights are chosen as

$$\omega_r = \frac{\alpha_r}{\sum_{s=0}^{m-1} \alpha_s}, \quad \alpha_r = \frac{d_r}{(\epsilon + \beta_r)^2}.$$

Here,  $d_r$  are the coefficients such that  $\sum_{r=0}^{m-1} d_r v_{i+1/2}^{(r)}$  coincides with the reconstruction using all the  $(2m - 1)$  cell averages. Show that

- (a) For  $m = 2$ ,  $d_0 = \frac{2}{3}$ ,  $d_1 = \frac{1}{3}$ ;
- (b) For  $m = 3$ ,  $d_0 = \frac{3}{10}$ ,  $d_1 = \frac{3}{5}$ ,  $d_2 = \frac{1}{10}$ ;
- (c)  $d_r$  is away positive.

6. In the WENO reconstruction, let the reconstruction polynomial on the stencil  $S_r(i)$  be denoted by  $p_r(x)$ . Define

$$\beta_r = \sum_{l=1}^{m-1} \int_{I_i} h^{2l-1} \left( \frac{\partial^l p_r(x)}{\partial x^l} \right) dx \quad r = 0, \dots, m-1.$$

- (a) Show that when  $m = 2$ ,

$$\beta_0 = (\bar{v}_{i+1} - \bar{v}_i)^2, \quad \beta_1 = (\bar{v}_i - \bar{v}_{i-1})^2.$$

- (b) Verify the condition that

$$\beta_r = D(1 + \mathcal{O}(h^{m-1})) \tag{1}$$

when  $m = 2$ , where  $D$  is a nonzero quantity independent of  $r$  (but may depend on  $h$ ).

- (c) Find the expression of  $\beta_0$ ,  $\beta_1$  and  $\beta_2$  for  $m = 3$  and verify the condition (1).

7. Consider the elliptic problem in non-divergence form:

$$\mathbf{A}(x) : D^2 u(x) = f(x), \quad \mathbf{A}(x) \geq 0.$$

If there exists  $\Lambda_0 \geq \lambda_0 > 0$ , such that  $\lambda_0 \mathbf{I} \leq \mathbf{A}(x) \leq \Lambda_0 \mathbf{I}$  for all  $x \in \Omega$ , show that the non-divergence form is uniformly elliptic.