## 微分拓扑, 2018 年春季

## 作业 2

## 上交时间及方式: 2017.04.10

**REQUIRED**:

- 1. Let X and Z be transversal submanifolds of Y. Show that if  $y \in X \cap Z$ , then  $T_y(X \cap Z) = T_y(X) \cap T_y(Z)$ , that is, the tangent space to the intersection is the intersection of the tangent spaces.
- 2. If  $f: M \to Y$  and  $g: N \to Y$  are smooth morphisms of manifolds, the fiber product  $M \times_Y N$  is defined by

$$M \times_Y N = \{ (m, n) \mid m \in M, n \in N, f(m) = g(n) \}$$

We say that f and g are transverse if for all pairs  $(m,n) \in M \times_Y N$  we have that  $Df_m(T_mM) + Dg_n(T_nN) = T_{f(m)}Y$ . Show that if  $g: N \to Y$  is an embedding of a submanifold, that f is transverse to g in this sense if and only if f is transverse to g(N). Show that if f and g are transverse, the fiber product  $M \times_Y N$  is a manifold.

(Hint: Consider the map  $f \times g : M \times N \to Y \times Y$ .)

3. If  $M \subset \mathbb{R}^n$  is a submanifold, define the tangent bundle to M to be the set

$$TM = \{(m, v) \mid m \in M, v \in T_mM\} \subset M \times R^n$$

Show that TM is a smooth manifold. Show that if  $f: M \to N$  is a smooth morphism, that the natural map  $Df: TM \to TN$  defined by  $Df(m, v) = (f(m), Df_m(v))$  is a smooth morphism. Conclude that if M is diffeomorphic to N, then TM is diffeomorphic to TN. (So that TM does not depend on how M was embedded into  $\mathbb{R}^n$ .)

- 4. Show that diffeomorphism is a stable property for maps of compact manifolds. That is, if  $f_0: X \to Y$  is a diffeomorphism between compact smooth manifolds, and  $f_t: X \to Y$  is a homotopy of  $f_0$ , then, for some  $\varepsilon > 0$ , each  $f_t$  with  $t < \varepsilon$  is also a diffeomorphism. Here we suppose X and Y are connected.
- 5. Suppose Z is a submanifold of X with  $\dim Z < \dim X$ . Show that Z has measure zero in X WITHOUT using Sard.
- 6. Let X be a compact manifold. Show that there exist Morse functions on X which take distinct values at distinct critical points.

(Hint: Let  $x_1, \dots, x_N$  be the list of critical points of f. Let  $\rho_i$  be a smooth function which is one on a small neighborhood of  $x_i$  and zero outside a slightly larger neighborhood. Choose numbers  $a_1, \dots, a_N$  such that

$$f(x_i) + a_i \neq f(x_j) + a_j$$
 if  $i \neq j$ 

Show that for  $a_i$  small enough, the function  $f + \sum a_i \rho_i$  is Morse and has the same critical points as f and is even arbitrarily close to f.)

- 7. A vector field  $\vec{v}$  on a manifold X in  $\mathbb{R}^N$  is a smooth map  $\vec{v} : X \to \mathbb{R}^N$  such that  $\vec{v}(x)$  is always tangent to X at x. Show that a vector field  $\vec{v}$  is a cross section of T(X) (that is, a smooth map  $\vec{v} : X \to T(X)$  such that  $p \circ \vec{v} = \mathbb{1}_X$ ), and vice versa. Here p is the projection map  $p : T(X) \to X$  defined by p(x, v) = x.
- 8. (*The Whitney Immersion Theorem*) Show that every k-dimensional manifold X may be immersed into  $\mathbb{R}^{2k}$ .
- 9. (*The Smooth Urysohn Theorem*) If A and B are disjoint, smooth, closed subsets of a manifold X, prove that there is a smooth function  $\phi$  on X such that  $0 \le \phi \le 1$  with  $\phi = 0$  on A and  $\phi = 1$  on B.

(Hint: Partition of Unity.)

## SUGGESTED:

- 1. \* If f and g are two smooth maps from a smooth manifold M to  $S^n$ , show that if for all  $m \in M$  one has that ||f(m) g(m)|| < 2 then f is smoothly homotopic to g.
- 2. \* Exhibit a smooth map  $f : \mathbb{R} \to \mathbb{R}$  whose set of critical values is dense.

(Hint: Write the rationals in a sequence  $r_0, r_1, \cdots$ . Now construct a smooth function on [i, i + 1] that is zero near the endpoints and that has  $r_i$  as a critical value.)

- 3. \* Show that the sphere  $S^k$  is simply connected if k > 1 using Sard. (Hint: If  $f : S^1 \to S^k$  and k > 1, Sard gives you a point  $p \notin f(S^1)$ . Now use stereographic projection.)
- 4. \* Show that  $T(X \times Y)$  is diffeomorphic to  $T(X) \times T(Y)$ , and the projection  $p: T(X) \to X$ , p(x, v) = x is a submersion.
- 5. \* Let S(X) be the set of points  $(x, v) \in T(X)$  with |v| = 1. Show that S(X) is a smooth submanifold of T(X) of dimension 2k 1. S(X) is called the *unit tangent bundle* or *sphere tangent bundle* of X.

(Hint: Consider the map  $(x, v) \mapsto |v|$ .)